

บทที่ 6
เอกสารอ้างอิง

1. Milne, J.A., *The metabolism of androgens by sebaceous glands*. Br J Dermatol, 1969. 81: p. Suppl 2:23-8.
2. Birgham, K.D. and D.A. Shaw, *The metabolism of testosterone by human male scalp skin*. J Endocrinol, 1973. 57: p. 111-21.
3. Liang, T., G.H. Rasmusson, and J.R. Brooks, *12. Androgens: Pharmacodynamics and antagonists. Biochemical and biological studies with 4-aza-steroidal 5 alpha-reductase inhibitors*. J Steroid Biochem, 1983. 19(1A): p. 385-90.
4. Mauvais-Jarvis, P., J.P. Bercovici, and F. Gauthier, *In vivo studies on testosterone metabolism by skin of normal males and patients with the syndrome of testicular feminization*. J Clin Endocrinol Metab, 1969. 29(3): p. 417-21.
5. Wilson, J.D. and J.D. Walker, *The conversion of testosterone to 5 alpha-androstan-17 beta-ol-3-one (dihydrotestosterone) by skin slices of man*. J Clin Invest, 1969. 48(2): p. 371-9.
6. Labrie, F., *Intracrinology*. Mol Cell Endocrinol, 1991. 78(3): p. C113-8.
7. Chen, W., C.C. Zouboulis, and C.E. Orfanos, *The 5 alpha-reductase system and its inhibitors. Recent development and its perspective in treating androgen-dependent skin disorders*. Dermatology, 1996. 193(3): p. 177-84.
8. Matias, J.R., V.L. Malloy, and N. Orentreich, *Synergistic antiandrogenic effects of topical combinations of 5 alpha-reductase and androgen receptor inhibitors in the hamster sebaceous glands*. J Invest Dermatol, 1988. 91(5): p. 429-33.
9. Burton, J.L., U. Laschet, and S. Shuster, *Reduction of sebum excretion in man by the antiandrogen, cyproterone acetate*. Br J Dermatol, 1973. 89(5): p. 487-90.
10. Matias, J.R. and M. Gaillard, *Local inhibition of sebaceous gland growth by topically applied RU 58841*. Ann N Y Acad Sci, 1995. 761: p. 56-65.
11. Matias, J.R., et al., *The lack of effect of 11 alpha-hydroxyprogesterone on the flank-organ and ear sebaceous glands of adult male Syrian golden hamsters*. Arch Dermatol Res, 1984. 276(5): p. 346-8.
12. Ye, F., et al., *Effects of topical antiandrogen and 5-alpha-reductase inhibitors on sebaceous glands in male fuzzy rats*. Skin Pharmacol, 1997. 10(5-6): p. 288-97.

13. Brooks, J.R., et al., *Topical anti-androgenicity of a new 4-azasteroid in the hamster*. Steroids, 1991. 56(8): p. 428-33.
14. Luderschmidt, C., et al., *17 alpha-Propylmesterolone (SH 434): an antiandrogenic sebosuppressive substance not influencing circulating testosterone concentrations. Experimental studies in Syrian hamsters*. Naunyn Schmiedebergs Arch Pharmacol, 1984. 328(2): p. 214-8.
15. Chansakaow, S., et al., *Isoflavonoids from Pueraria mirifica and their estrogenic activity*. Planta Med, 2000. 66(6): p. 572-5.
16. Cain, J.C., *Miroestrol: an oestrogenic from the plant Pueraria mirifica*. Nature, 1960. 188: p. 774-7.
17. Kulling, S.E. and M. Metzler, *Induction of micronuclei, DNA strand breaks and HPRT mutations in cultured Chinese hamster V79 cells by the phytoestrogen coumestrol*. Food Chem Toxicol, 1997. 35(6): p. 605-13.
18. Whitten, P.L., et al., *Phytoestrogen influences on the development of behavior and gonadotropin function*. Proc Soc Exp Biol Med, 1995. 208(1): p. 82-6.
19. Whitten, P.L., et al., *Potential adverse effects of phytoestrogens*. J Nutr, 1995. 125(3 Suppl): p. 771S-776S.
20. Kashemsanta, L., K. Suvatabandhu, and A.K. Airy Shaw, *A new species of Pueraria (Leguminosae) from Thailand, yielding an oestrogenic principle*. Kew Bull, 1952. 4: p. 549-51.
21. van der Maesen, L.J.T., *Revision of the genus Pueraria DC. with some note on Teyleria Backer*. Agric Univ Wageningen Papers, 1985. 85(1): p. 5-132.
22. Chen, C., et al., *Local and systemic reduction by topical finasteride or flutamide of hamster flank organ size and enzyme activity*. J Invest Dermatol, 1995. 105(5): p. 678-82.