บรรณานุกรม

พิชัย คงพิทักษ์ การใช้ผึ้งพันธุ์ (*Apis mellifera* L.) เป็นแมลงผสมเกสรในการผลิตเมล็ดทานตะวัน (*Helianthuse annuus* L.) วิทยานิพนธ์ มหาวิทยาลัยเกษตรศาสตร์ 2530.

Abrams, R.L., C.R. Edwards and T. Harris, 1978. Yields and cross-polination of soybean as affected by honeybees and alfalfa leafcutting bees. Am. Bee J. 118 : 555-558.

Beard, B.H. and P.F. Knowles, 1971. Frequency of cross-polination of soybeans after irradiation. Crop Sci 11 : 489 - 492.

Boerma, H.R., and A. Moradshahi, 1975. Pollen movement within and between rows to male - sterile within and between rows to male - sterile soybean. Crop Sci 15 : 858 - 861.

Caviness, C.E., 1966. Estimates of natural cross - pollination in Jackson soybean in Arkansas. Crop Sci 6 : 211 - 213.

Cutter, C.H., 1934. A Simple method for making soybean hybrids. J. Am. Soc. Agron. 26 : 252 - 254.

Erickson, E.H., G.A. Berger, J.G. Shannon and J.M. Robins, 1978. Honeybee pollination incresed soybean yields in the Missisipppi delta region of Arkansas and Missouri. J. Econ. Entomol. 71 : 601-603.