



บรรณานุกรม

- จุรีพร จิตจำรูญโขคไชย. (2537). ผลของการออกกำลังกายแบบใช้ออกซิเจนต่อระดับไขมันในเลือด. *รามาธิบดีเวชสาร*, 17(1), 20-26.
- เพ็ญพิมล รั้มมรรคคิต. (2537). สรีรวิทยาของการออกกำลังกาย. ในวิรุฬห์ เหล่าภารเกษม (บรรณาธิการ), *กีฬาเวชศาสตร์* (หน้า 15-18). กรุงเทพฯ: พ.บ.ฟอร์ม บุคส์ เซ็นเตอร์.
- วิศาล คันธารัตนกุล. (2549). Physical fitness testing:Clinical implication. ใน กุลวีณ์ สุจิริตการ (บรรณาธิการ), *การอบรมเชิงปฏิบัติการสรีรวิทยา-พยาธิสรีรวิทยา* ครั้งที่ 24. (หน้า 60-61). กรุงเทพฯ: คุลเซอร์ฟ.
- สมชาย จักรพันธ์, ละอียด ชูประยูร, วนชัย ไชยศิทธิ์, สุพิน พรพิพัฒน์กุล, สุจิตรา อุสาหะ, จำปา พ่องเงิน และคณะ. (2542). รายงานวิจัยการพัฒนาแบบประเมินและวิเคราะห์ความเครียดด้วยตนเองสำหรับประชาชนไทยด้วยคอมพิวเตอร์ (พิมพ์ครั้งที่ 2). สมุทรปราการ: ทีคอม.
- สมชาย ลีทองอิน. (2542). มุมมองใหม่ในการออกกำลังกายเพื่อสุขภาพ. *สถานการณ์ด้านการออกกำลังกาย*, 4 (5), 12.
- สมบัติ ดาปัญญา. (2526). คู่มือคลายเครียด ความรู้เรื่องความเครียด การป้องกันและแก้ไข (พิมพ์ครั้งที่ 2) กรุงเทพฯ: เนริยณบุญการพิมพ์
- สำนักงานสถิติแห่งชาติ. (ม.บ.บ.) การสำรวจพฤติกรรมการสูบบุหรี่และการดื่มสุราของประชากร พ.ศ.2547. สืบค้นเมื่อ 4 ธันวาคม 2551,
จาก http://service.nso.go.th/nso/nsopublish/service/survey/cigar_drink_47.pdf
- เสก อักษรานุเคราะห์. (2530). พลังงานสำหรับการออกกำลังกาย. *จุฬาลงกรณ์เวชสาร*, 31 (4), 271-286.
- Ainsworth, B.E. Haskell, W.L., Whitt, M.C., Irwin, M.L., Swartz, A.M., Strath, S.J., et al. (1993). Compendium of Physical Activities: Classification of energy cost of human physical activities. *Medicine and Science in Sports and Exercise*, 25, 71-80.
- Alberti, K.G.M.M, Zimmet, P.Z., and Shaw, J.E. (2005). The metabolic syndrome: a new world-wide definition from the International Diabetes Federation consensus. *Lancet*, 366 (9491), 1059-1062.

- Alvin, C. (2004). Keeping Metabolic Syndrome in Check. Retrieved August 9, 2006, from <http://healthlink.mcw.edu/article/1031002423.html>.
- Anderson, E.A., Hoffman, R. P., Balon, T. W., Sinkey, C. A., and Mark, A. L. (1991). Hyperinsulinemia produces both sympathetic neural activation and vasodilation in normal humans. *The Journal of Clinical Investigation*, 87(6), 2246-2252.
- Attvall, S. Fowelin, J., Lager, I., Schenck, H.V., and Smith, U. (1993). Smoking induces insulin resistance-a potential link with the insulin resistance syndrome. *Journal of Internal Medicine*, 233, 327-332.
- Bahia, L., Aguiar, L.G., Villela, N., Bottino, D., and Godoy, A.F. (2006). Relationship between adipokines, inflammation, and vascular reactivity in lean controls and obese subjects with metabolic syndrome. *Clinics*, 61(5), 433-440.
- Bamia, C., Trichopoulou, A., Lenas, D., and Trichopoulos, D. (2004). Tobacco smoking in relation to body fat mass and distribution in a general population sample. *International Journal of Obesity and Related Metabolic Disorder*, 28(8), 1091-1096.
- Bjorntorp, P. (1991). Visceral fat accumulation-the missing link between psychosocial factors and cardiovascular disease. *Journal of Internal Medicine*, 230, 195-201.
- Blair, S.N., Kohl, H.W., Barlow, C.E., Paffenbarger, R.S., Gibbons, L.W., and Macera, C.A. (1995). Changes in physical fitness and all-cause mortality: a prospective study of healthy and unhealthy men. *The Journal of The American Medical Association*, 273, 1093-1098.
- Blair, S.N., Cheng, Y., and Holder, J.S. (2001). Is physical activity or physical fitness more important in defining health benefits? *Medicine and Science in Sports and Exercise*, 33(6), S379-S399.
- Bloom, Z.T. (2002). Obesity, hypertension, and insulin resistance. *Diabetes Care*, 25, 2088-2097.
- Bonadonna, R.C. (2006). The Metabolic Syndrome Is a Risk Indicator of Microvascular and Macrovascular Complications in Diabetes. *Diabetes Care*, 29, 2701-2707.

- Brien, S.E., and Katzmarzyk, P.T. (2006). Physical activity and the metabolic syndrome in Canada. *Applied Physiology, Nutrition, and Metabolism*, 31, 40-47.
- Brunner, E.J., Marmot, M.G., Nanchahal, K., Shipley, M.J., Stansfeld, S.A., Juneja, M., et al. (1997). Social inequality in coronary risk: central obesity and the metabolic syndrome. Evidence from the Whitehall II study. *Diabetologia*, 40, 1341–1349.
- Brunner, E.J., Hemingway, H., Walker, B.R., Page, M., Clarke, P., Juneja, M., et al. (2002). Adrenocortical, autonomic, and inflammatory causes of the metabolic syndrome-nested case-control study. *Circulation*, 106, 2659-2665.
- Carnethon, M.R., Loria, C.M., Hill, J.D., Sidney, S., and Savage, P.J. (2004). Risk factors for the metabolic syndrome. *Diabetes Care*, 27, 2707-2715.
- Carroll, S., and Dudfield, M. (2004). What is the relationship between exercise and metabolic abnormalities? A review of the metabolic syndrome. *Sports Medicine*, 34(6), 371-418.
- Chandola, T., Brunner, E., and Marmot, M. (2006). Chronic stress at work and the metabolic syndrome: prospective study. *British Medical Journal*, 332, 521-525.
- Choiniere, R., Lafontaine, P., and Edwards, A.C. (2000). Distribution of cardiovascular disease risk factors by socioeconomic status among Canadian adults. *Canadian Medical Association Journal*, 162, S13-S24.
- de Simone, G., Devereux, R.B., Chinali, M., Best, L.G., Lee, E.T., and Galloway, J.M. (2007). Prognostic impact of metabolic syndrome by different definitions in a population with high prevalence of obesity and diabetes: the Strong Heart Study. *Diabetes Care*, 30(7), 1851-1856.
- Deen, D. (2004). Metabolic Syndrome: Time for Action. *American Family Physician*, 69(12), 2875-2882.
- Einhorn, D., Reaven, G.M., Cobin, R.H., Ford, E., Ganda, O.P., Handelsman, Y., et al. (2003). American College of Endocrinology position statement on the insulin resistance syndrome. *Endocrine Practice*, 9(3), 237-252.

- Farmer, A. (2006). Metabolic syndrome and mortality. *British Medical Journal*, 332 (7546), 882.
- Fernando, G.R., and Martha, Rodr. (2005). Concordance Between the 2005 International Diabetes Federation Definition for Diagnosing Metabolic Syndrome With the National Cholesterol Education Program Adult Treatment Panel III and the World Health Organization Definitions. *Diabetes Care*, 28, 2588-2589.
- Folkow, B. (1997). Physiological aspects of the "defense" and "defeat" reactions. *Acta Physiologica Scandinavica*, 161 (640), 34-37.
- Ford, E.S., and Giles, W.H. (2002). Serum C-reactive protein and fibrinogen concentrations and self-reported angina pectoris and myocardial infarction. *Arteriosclerosis, thrombosis, and vascular biology*, 20, 1052-1056.
- Ford, E.S., and Giles, W.H., (2003). A Comparison of the Prevalence of the Metabolic Syndrome Using Two Proposed Definitions. *Diabetes Care*, 26, 575-581.
- Ford, E.S., Giles, W.H., and Dietz, W.H. (2002). Prevalence of the metabolic syndrome among US adults: findings from the Third National Health and Nutrition Examination Survey. *The Journal of The American Medical Association*, 287, 356-359.
- Franklin, B.A., Swain, D.P., and Shephard, R.J. (2003). New insights in the prescription of exercise for coronary patients. *The Journal of Cardiovascular Nursing*, 18, 116-123.
- Giri, S., Thompson, P.D., Kiernan, F.J., Clive, J., Daniel B. Fram, D.B., Mitchel, J.F., et al. (1999). Clinical and angiographic characteristics of exertion-related acute myocardial infarction. *The Journal of The American Medical Association*, 282, 1731-1736.
- Glanz, K., Lewis, FM. and Rimer, B.K. (Eds.) (1997). *Health behavior and health education*. San Francisco: Jossey-Bass.

- Godsland, I.F., Leyva, F., Walton, C., Worthington, M., and Stevenson, J.C. (1998). Associations of smoking, alcohol and physical activity with risk factors for coronary heart disease and diabetes in the first follow-up cohort of the Heart Disease and Diabetes Risk Indicators in a Screened Cohort study (HDDRISC-1). *Journal of Internal Medicine*, 244(1), 33-41.
- Goodpaster, B.H., He, J., Watkins, S., and Kelley, D.E. (2001). Skeletal muscle lipid content and insulin resistance: evidence for a paradox in endurance-trained athletes. *Journal of Clinical Endocrinology & Metabolism*, 86, 5755-5761.
- Grundy, S.M. (2002). Obesity, metabolic syndrome, and coronary atherosclerosis. *Circulation*, 105, 2696-2698
- Grundy, S.M. (2006). Does the Metabolic Syndrome Exist?. *Diabetes Care*, 29(7), 1689-1692.
- Grundy, S.M., Brewer, B.H., Cleeman, J.L., Smith, S.C., and Lenfant, C. (2004). Definition of Metabolic Syndrome. *Circulation*, 109, 433-438.
- Grundy, S.M., Cleeman, J.L., Daniels, S.R., Donato, K.A., Eckel, R.H., Barry, A., et al. (2005). Diagnosis and Management of the Metabolic Syndrome.: a statement for health care professionals: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation*, 112, 2735-2752.
- Haffner, S.M., Valdez, A., Hazudah, P., Mitchellb, D., Morales, P.A., Stern, M.P., et al. (1992). Prospective analysis of the insulin-resistance syndrome (syndrome X). *Diabetes*, 41(6), 715-722.
- Hagberg, J.M., Montain, S.J., Martin, W.H., and Ehsani, A.A. (1989). Effect of exercise training in 60 to 69 years old persons with essential hypertension. *The American Journal of Cardiology*, 64(5), 348-353.
- Hagberg, J.M., Park, J.J., and Brown, M.D. (2000). The role of exercise training in the treatment of hypertension. *Sports Medicine*, 30, 193-206.
- Haines, C. (2005). Weight Loss: What Is Obesity? Retrieved July 25, 2006, from http://www.webmd.com/content/article/46/2731_1678.htm

- Halbert, J.A., Silagy, C.A., Finucane, P., Withers, R.T., Hamdorf, P.A., and Andrews, G.R. (1997). The effectiveness of exercise training in lowering blood pressure: a meta-analysis of randomized controlled trials of 4 weeks or longer. *Journal of Human Hypertension*, 11(10), 641-649.
- Hambrecht , R., Niebauer, J., Marburger, C., Grunze, M., Kalberer, B., Hauer, K., Schlierf, G., et al. (1993). Various intensities of leisure time physical activity in patients with coronary artery disease: effects on cardiorespiratory fitness and progression of coronary atherosclerotic lesions. *Journal of the American College of Cardiology*, 22, 468-477.
- Heim, D.L, Holcomb, C.A., and Loughin, T.M. (2000). Exercise mitigates the association of abdominal obesity with high-density lipoprotein cholesterol in premenopausal women: results from the third National Health and Nutrition Examination Survey. *Journal of the American Dietetic Association*, 100, 1347-1353.
- Hinderliter, A., Sherwood, A., Gullette, E.C.D., Babyak, M., Waugh, R., Georgiades, A., et al. (2002). Reduction of left ventricular hypertrophy after exercise and weight loss in overweight patients with mild hypertension. *Archives of Internal Medicine*, 162, 1333-1339.
- Hu, F.B., Stampfer, M.J., Solomon, C., Liu, S., Colditz, G.A., Speizer, F.E., et al (2001). Physical activity and risk for cardiovascular events in diabetic women. *Annals of Internal Medicine*, 134(2), 96-105.
- Hu, F.B., Willett, W.C., Li, T., Stampfer, M.J., Colditz, G.A., and Manson, J.E. (2004). Adiposity as compared with physical activity in predicting mortality among women. *The New England Journal of Medicine*, 351, 2694-2703.
- Hu, G., Qiao,Q., Tuomilehto, J., Balkau, B., Borch, J.K., and Pyorala, K. (2004). Prevalence of the metabolic syndrome and its relation to all-cause and cardiovascular mortality in nondiabetic European men and women. *Archives of Internal Medicine*, 164(10), 1066-1076.

- Hunt, K.J., Resendez, R.G., Williams, K., Haffner, S.M., and Stern, M.P. (2004). National Cholesterol Education Program versus World Health Organization metabolic syndrome in relation to all-cause and cardiovascular mortality in the San Antonio Heart Study. *Circulation*, 110, 1251–1257.
- Isomaa, B., Almgren, P., Tuomi, T., Forsen, G., Lahti, K., and Nissen, M. (2001). Cardiovascular morbidity and mortality associated with the metabolic syndrome. *Diabetes Care*, 24, 683–689.
- Jatuporn, S., Sangwananaroj, S., and Saengsiri, A. (2003). Short-term effects of an intensive lifestyle modification program on lipid peroxidation and antioxidant systems in patients with coronary artery disease. *Clinical Hemorheology and Microcirculation*, 29, 429-436.
- Jequier, E. (2002). Leptin signaling, adiposity, and energy balance. *Annals of the New York Academy of Sciences*, 967, 379-388.
- Johnson, J.L., Slentz, C.A., Houmard, J.A., Samsa, G.P., Duscha, B.D., Aiken, L.B., et al. (2007). Exercise Training Amount and Intensity Effects on Metabolic Syndrome (from Studies of a Targeted Risk Reduction Intervention through Defined Exercise). *The American Journal of Cardiology*, 100(12), 1759-1766.
- Jurca, R. Lamonte, M.J., Barlow, C.E., Kampert, J.B., Church, T.S., and Blair ,S.N. (2005). Association of Muscular Strength with Incidence of Metabolic Syndrome in Men. *Medicine&Science in Sports & Exercise*, 37(11), 1849-1855.
- Kahn, R., Buse, J., Ferrannini, E., and Stern, M. (2005). The metabolic syndrome: time for a critical appraisal: joint statement from the American Diabetes Association and the European Association for the Study of Diabetes. *Diabetes Care*, 28, 2289–2304.
- Kaplan, N.M. (1989). The deadly quartet. Upper-body obesity, glucose intolerance, hypertriglyceridemia, and hypertension. *Archives of Internal Medicine*, 149 (7), 1514–1520
- Kaplan, G.A., and Keil J.E. (1993). Socioeconomic factors and cardiovascular disease: a review of the literature. *Circulation*, 88, 1973–1998.

- Kaplan, G.A., Strawbridge, W.J., Cohen, R.D., and Hungerford, L.R. (1996). Natural history of leisure-time physical activity and its correlates: Associations with mortality from all causes and cardiovascular diseases over 28 years. *American Journal of Epidemiology*, 144(8), 793-797..
- Kasai, T., Miyauchi, K., Kubota, N., Tamura, H., Kojima, T., Yokoyama, K., et al. (2008) The relationship between the metabolic syndrome defined by various criteria and the extent of coronary artery disease. *Atherosclerosis*, 197(2), 944-950.
- Kim, Y.H., and Yang, Y.O. (2005). Effects of walking exercise on metabolic syndrome risk factors and body composition in obese middle school girls. *Taehan Kanho Hakhoe Chi*, 35(5), 858-867.
- Kodama, S., Tanaka, S., Saito, K., Shu, M., Sone, Y., Onitake, F., et al. (2007). Effect of Aerobic Exercise Training on Serum Levels of High-Density Lipoprotein Cholesterol. *Archives of Internal Medicine*, 167(10), 999-1008.
- Koeslag, J.H., Saunders, P.T., and Terblanche, E. (2003). A reappraisal of the blood glucose homeostat which comprehensively explains the type 2 diabetes mellitus-syndrome X complex. *Journal of Physiology*, 549, 333-346.
- Kong, C. (2001). Smoking is associated with increased hepatic lipase activity, insulin resistance, dyslipidaemia and early atherosclerosis in Type 2 diabetes. *Atherosclerosis*, 156, 373-78.
- Kraja, A.T., Borecki, I.B., North, K., Tang, W., Myers, R.H., and Hopkins, P.N. (2006). Longitudinal and age trends of metabolic syndrome and its risk factors: The Family Heart Study. *Nutrition&Metabolism*, 3, 41.
- Kuusisto, J., Lempiäinen, P., Mykkänen, L., and Laakso, M. (2001). Insulin resistance syndrome predicts coronary heart disease events in elderly type 2 diabetic men. *Diabetes Care*, 24, 1629-1633.

- Laaksonen, D.E., Lakka, H.M., Niskanen, L.K., Kaplan, G.A., Salonen, J.T., and Lakka T.A. (2002). Metabolic syndrome and development of diabetes mellitus: application and validation of recently suggested definitions of the metabolic syndrome in a prospective cohort study. *American Journal of Epidemiology*, 156, 1070-1077.
- Laaksonen, D.E., Lindström, J., Lakka, T.A., Eriksson, J.G., Niskanen, L.K., Wikström, K., et al. (2005). Physical activity in the prevention of type 2 diabetes. *Diabetes*, 54, 158-165.
- Lakka, H.M., Laaksonen, D.E., Lakka, T.A., Niskanen, L.K., Kumpusalo, E., Tuomilehto, J., et al. (2002). The metabolic syndrome and total and cardiovascular disease mortality in middle-aged men. *The Journal of The American Medical Association*, 288, 2709-2716.
- Langin, D. (2001). Diabetes, insulin secretion, and the pancreatic beta-cell mitochondrion. *The New England Journal of Medicine*, 345, 1772-1774.
- Lee, I.M., Hsieh, C.C., and Paffenbarger, R.S. (1995). Exercise intensity and longevity in men: the Harvard Alumni Health Study. *The Journal of The American Medical Association*, 273, 1179-1184.
- Lee, K.W., and Lip, G.Y.H. (2003). Effects of lifestyle on hemostasis, fibrinolysis, and platelet reactivity: a systematic review. *Archives of Internal Medicine*, 163, 2368-2392.
- Lee, K.W., and Lip, G.Y.H. (2004). Acute versus habitual exercise, thrombogenesis and exercise intensity. *Thrombosis and Haemostasis*, 91, 416-419.
- Legro, R.S. (2001). Diabetes prevalence and risk factors in polycystic ovary syndrome. *Obstetrics and Gynecology Clinics of North America*, 28(1), 99-109.
- Leon, A.S., Connett, J., Jacobs, D.R., and Rauramaa, R. (1987). Leisure-time physical activity levels and risk of coronary heart disease and death. The Multiple Risk Factor Intervention Trial. *The Journal of the American Medical Association*, 258 (17), 2388-2395.

- Lempainen, P., Mykkänen, L., Pyörälä, K., Laakso, M., and Kuusisto, J. (1999). Insulin resistance syndrome predicts coronary heart disease events in elderly non-diabetic men. *Circulation*, 100, 123–128.
- Liu, S., and Manson, J.E. (2001). Dietary carbohydrates, physical inactivity, obesity, and the 'metabolic syndrome' as predictors of coronary heart disease. *Current Opinion in Lipidology*, 12, 395–404.
- Lohsoontorn, V., Bodi, D., and Michelle, A.W. (2006). Prevalence of metabolic syndrome and its relationship to white blood cell count in population of Thai men and women receiving routine health examination. *American Journal of Hypertension*, 19, 339-345.
- Loreto, C.D., Fanelli, C., Lucidi, P., Murdolo, G., Cicco, A.D., Parlanti, N., et al. (2005) Long-term impact of different amounts of physical activity on type 2 diabetes *Diabetes Care*, 28, 1295-1302.
- Maggi, S., Noale, M., Gallina, P., Bianchi, D., Marzari, C., Limongi, F., et al. (2006). Metabolic syndrome, diabetes, and cardiovascular disease in an elderly Caucasian cohort: the Italian Longitudinal Study on Aging. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 61(5), 505-510.
- Malhotra, V., Singh, S., Tandon, O.P., and Sharma, S.B. (2005). The beneficial effect of yoga in diabetes. *Nepal Medical Collage Journal*, 7(2), 145-147.
- Masulli, M., and Vaccaro, O. (2006). Association Between Cigarette Smoking and Metabolic Syndrome. *Diabetes Care*, 29, 482-483.
- McArdle, W.D., Katch, F.I., and Katch, V.L. (2001). *Exercise Physiology: Energy, Nutrition and Human Performance*. U.S.A: Lippincott.Williams&Wilkins.
- McKeigue, P.M., Shah, B., and Marmot, M.G. (1991). Relation of central obesity and insulin resistance with high diabetes prevalence and cardiovascular risk in South Asians. *Lancet*, 337, 382–86.

- Michael, J.L. (2005). Cardiorespiratory Fitness Is Inversely Associated With the Incidence of Metabolic Syndrome. *Circulation*, 112, 505-512.
- Mittleman, M.A., Maclure, M., Tofler, G.H., Sherwood, J.B., Goldberg, R.J., and Muller, J.E. (1993). Triggering of acute myocardial infarction by heavy physical exertion. Protection against triggering by regular exertion. *The New England Journal of Medicine*, 329(23), 1677-1683.
- Mohan, V., Gokulakrishnan, K., Deepa, R., Shanthirani, C.S., and Datta, M. (2005). Association of physical inactivity with components of metabolic syndrome and coronary artery disease--the Chennai Urban Population Study (CUPS no. 15). *Diabetic Medicine*, 22(9), 1206-1211.
- Montague, C.T., and O'Rahilly, S. (2000). The perils of portliness: Causes and consequences of visceral obesity. *Diabetes*, 49, 883-888.
- Onat, A., Ceyhan, K., Basar, O., Erer, B., Toprak, S., and Sansoy, V. (2002). Metabolic syndrome: major impact on coronary risk in a population with low cholesterol levels: a prospective and cross-sectional evaluation. *Atherosclerosis*, 165, 285-292.
- Orio, F., Giallauria, F., Palomba, S., Cascella, T., Manguso, F., Vuolo, L., et al. (2006). Cardiopulmonary Impairment in Young Women with Polycystic Ovary Syndrome. *The Journal of Clinical Endocrinology & Metabolism*, 91(8), 2967-2971.
- Paffenbarger, R.S., Jung, D.L., Leung, R.W., and Hyde, R.T. (1991). Physical activity and hypertension: an epidemiological view. *Annals of Medicine*, 23, 319-327.
- Paffenbarger, R.S., Hyde, R.T., Wing, A.L., Lee, I.M., Jung, D.L., and Kampert, J.B. (1993). The association of changes in physical-activity level and other lifestyle characteristics with mortality among men. *The New England Journal of Medicine*, 328(8), 574-576.
- Paffenbarger, R.S., Kampert J.B., Lee, I., Hyde, R.T., Leung, R.W., and Wing, A.L. (1994). Changes in physical activity and other lifeway patterns influencing longevity. *Medicine and Science in Sports and Exercise*, 26, 857-865.

- Park, H.S., Oh, S.W., Cho, S., Choi , W.H., and Kim, Y.S. (2004). The metabolic syndrome and associated lifestyle factors among South Korean adults. *International Journal of Epidemiology*, 33(2), 328-336.
- Park, T.G., Hong, H.R., Lee, J., and Kang, H.S. (2007). Lifestyle plus Exercise Intervention Improves Metabolic Syndrome Markers without Change in Adiponectin in Obese Girls. *Annals of Nutrition & Metabolism*, 51, 197-203.
- Park, Y.W., Zhu, S., Palaniappan, L., Heshka, S. Carnethon, M.R., and Heymsfield, S.B. (2003). The metabolic syndrome. Prevalence and associated risk factor findings in the US population from the Third National Helath and Nutrition Examination Survey. *Archives of Internal Medicine*, 163, 427-436.
- Patel, C., Marmot, M.G., and Terry, D.J. (1981). Controlled trial of biofeedback-aided behavioural methods in reducing mild hypertension. *British Medical Journal*, 282, 2005-2008.
- Patel, D.A., Srinivasan, S.R., Xu, J.H., Chen, W., and Berenson, G.S. (2006). Adiponectin and its correlates of cardiovascular risk in young adults: the Bogalusa Heart Study. *Metabolism* , 55(11),1551-1557.
- Pate, R.R., Pratt , M., Blair; S. N., Haskell, W. L., Macera, C. A., Bouchard, C., et al. (1995). Physical activity and public health: a recommendation from the Centers for Disease Control and Prevention and the American College of Sports Medicine. *The Journal of the American Medical Association*, 273, 402-407.
- Pearson, T.A. (2002). Consensus panel guide to comprehensive risk reduction for adult patients without coronary or other atherosclerotic vascular disease. *Circulation*, 99, 2480-2484.
- Pekkanen, J., Tuomilehto, J., Uutela, A., Vartiainen, E., and Nissinen, E. (1995). Social class, health behaviour, and mortality among men and women in eastern Finland. *British medical journal*, 311, 589-593.

- Pereria, M.A., FitzerGerald, S.J., Gregg, E.W., Joswiak, M.L., Ryan, W.J., Suminski, R.R., et al. (1997). A collection of physical activity questionnaires for health-related research. *Medicine and Science in Sports and Exercise*, 29, S146-S152.
- Pescatello, L.S., Murphy, D., and Costanzo, D. (2000). Low-intensity physical activity benefits blood lipids and lipoproteins in older adults living at home. *Age Ageing*, 29, 433-439.
- Phillips, D.I.W., Barker, D.J.P., Fall, C.H.D., Seckl, J.R., Whorwood, C.B., Wood, P.J., et al. (1998). Elevated plasma cortisol concentrations: a link between low birth weight and the insulin resistance syndrome. *The Journal of Clinical Endocrinology and Metabolism*, 83, 757-760.
- Pratley, R.E. (2000). Aerobic exercise training-induced reductions in abdominal fat and glucose-stimulated insulin responses in middle-aged and older men. *Journal of the American Geriatrics Society*, 48, 1055-1061.
- Purnell, J.Q., Kahn, S.E., Albers, J.J., Nevin, D.N., Brunzell, J.D., and Schwartz, R.S. (2000). Effect of weight loss with reduction of intraabdominal fat on lipid metabolism in older men. *The Journal of Clinical Endocrinology and Metabolism*, 85, 977-982.
- Pyorala, M., Pyörälä, M., Miettinen, H., Halonen, P., Laakso, M., and Pyörälä, K. (2000). Insulin resistance syndrome predicts the risk of coronary heart disease and stroke in healthy middle-aged men: the 22-year follow-up results of the Helsinki Policemen Study. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 20, 538-544.
- Rennie, K.L., McCarthy, N., Yazdgerdi, S., Marmot, M., and Brunner, E. (2003). Association of the metabolic syndrome with both vigorous and moderate physical activity. *International Journal of Epidemiology*, 32, 600-606.
- Resnik, H.E., and Howard, B.V. (2002). Diabetes and cardiovascular disease. *Annual Review of Medicine*, 53, 245-267.

- Roberts, C.K., Vaziri, N.D., and Barnard, R.J. (2002). Effect of diet and exercise intervention on blood pressure, insulin, oxidative stress, and nitric oxide availability. *Circulation*, 106, 2530-2532.
- Ross, R., Dagnone, D., Jones, P.J.H., Smith, H., Paddags, A., Hudson, R., et al. (2000). Reduction in obesity and related comorbid conditions after diet-induced weight loss or exercise-induced weight loss in men. A randomized, controlled trial. *Annals of Internal Medicine*, 133, 92-103.
- Rubin, D.B. (1987). *Multiple imputation for non-response in surveys*. New York: John Wiley.
- Salmenniemi, U., Ruotsalainen, E., Pihlajamäki, J., Vauhkonen, I., Kainulainen, S., Punnonen, K., et al. (2004). Multiple abnormalities in glucose and energy metabolism and coordinated changes in levels of adiponectin, cytokines, and adhesion molecules in subjects with metabolic syndrome. *Circulation*, 110(25), 3842-3848.
- Santos, A.C. (2005). Physical activity decreases the prevalence of metabolic syndrome. An observational study in Portugal American Public Health Association.
- Schooling, C.M., Thomas, G.N., Leung, G.M., Ho, S.Y., Janus, E.D. and Lam, T.H. (2007) Is Height Associated With Cardiovascular Risk in Chinese Adults? *Epidemiology*, 18(2), 274-278.
- Schwartz, R.S., Cain, K.C., Shuman, W. P., Larson, V., Stratton, J.R., Beard, J.C., Kahn, S.E., et al. (1992). Effect of intensive endurance training on lipoprotein profiles in young and older men. *Metabolism*, 41, 649-654.
- Scuteri, A., Najjar, S.S., Morrell, C.H., and Lakatta, E.G. (2005). The metabolic syncrome in older individuals: prevalence and prediction of cardiovascular events. *Diabetes Care*, 28, 882-887.
- Scuteri, A., Orru, M., Najjar, S.S., Albai, G., Lai, M., Cao, A., et al. (2006). Age-Associated Effects of the Metabolic Syndrome on Arterial Structure and Function. The ProgeNIA Study. *Circulation*, 114, 893-894.

- Shand, B.I., Scott, R.S., Elder, P.A., and George, P.M. (2003). Plasma adiponectin in overweight, nondiabetic individuals with or without insulin resistance. *Diabetes, Obesity and Metabolism*, 5(5), 349-353.
- Sheehan, M.T., and Jensen, M.D. (2000). Metabolic complications of obesity: Pathophysiologic considerations. *The Medical Clinics of North America*, 84, 373-385.
- Sherman, S.E., D'Agostino, R.B., Cobb, I.L., and Kannel, W.B. (1994). Physical activity and mortality in women in the Framingham Heart Study. *American Heart Journal*, 128, 879-884.
- Shimokata, H., Muller, D.C., and Andres, R. (1989). Studies in the distribution of body fat. III. Effects of cigarette smoking. *The Journal of The American Medical Association*, 261, 1169-1173.
- Shulman, G.I. (2000). Cellular mechanisms of insulin resistance. *The Journal of Clinical Investigation*, 107, 171-177.
- Siscovick, D.S., Weiss, N.S., Fletcher, R.H., and Lasky, T. (1984). The incidence of primary cardiac arrest during vigorous exercise. *The New England Journal of Medicine*, 311, 874-877.
- Skott, P., Vaag, A., Bruun, N. E., Hother-Nielsen, O., Gallet, M.A. , Beck-Nielsen, H., et. al. (1991). Effect of insulin on renal sodium handling in hyperinsulinaemic type 2 (non-insulin-dependent) diabetic patients with peripheral insulin resistance. *Diabetologia*, 34(4), 275-281.
- Stewart, K.J. (2002). Exercise training and the cardiovascular consequences of type 2 diabetes and hypertension: plausible mechanisms for improving cardiovascular health. *The Journal of The American Medical Association*, 288, 1622-1631.
- Straka, R.J. (2004). Pathophysiology: hypothesis of atherogenesis. Retrieved January 28, 2007, from <http://courses.ahc.umn.edu/pharmacy/5822/Atherogenesis.html>

- Sundström, J., Byberg, L., Zethelius, B., Lithell, H., and Lind, L. (2006). Clinical value of the metabolic syndrome for long term prediction of total and cardiovascular mortality: prospective, population based cohort study. *British Medical Journal*, 332(15), 878-882.
- Tabata, I. (2006). *Exercise and physical activity reference for health promotion 2006* (EPAR 2006): Physical activity, exercise, and physical fitness. (original work published in Japanese). Retrieved November 22, 2008, from <http://www.nih.go.jp/eiken/english/research/pdf/epar2006.pdf>
- Tanomsup, S., Aekplakom, W., Sritara, P., Woodward, M., Yamwong, S., Tunlayadechanont, S. et al. (2007). A comparison of components of two definitions of the metabolic syndrome related to cardiovascular disease and all-cause mortality in a cohort study in Thailand. *Diabetes Care*, 30(8), 2138-2140.
- Taylor, A.A. (2001). Pathophysiology of hypertension and endothelial dysfunction in patients with diabetes mellitus. *Endocrinology&Metabolism Clinics of North America*, 30, 983-997.
- Third report of the National Cholesterol Education Program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (Adult Treatment Panel III). Final report. (2002). *Circulation*, 106, 3143-3421.
- Thompson, P.D., Funk, E.J., Carleton, R. A., and Sturmer, W.Q. (1982). Incidence of death during jogging in Rhode Island from 1975 through 1980. *The Journal of The American Medical Association*, 247(18), 2535-2538.
- Trevisan, M., Liu, J., Bahsas, F.B., and Menotti, A. (1998). Syndrome X and mortality: A population-based study. Risk factor and life expectancy research group. *American Journal of Epidemiology*, 148(10), 958-966.
- Tricia, Y., Rana, J.S., Manson, J.E., Willett, W.C., Stampfer, M.J., Colditz, G.A., et al. (2006). Obesity as Compared With Physical Activity in Predicting Risk of Coronary Heart Disease in Women. *Circulation*, 113, 499-506.
- Vega, G.L. (2001). Obesity, the metabolic syndrome, and cardiovascular disease. *American Heart Journal*, 142, 1108-1116.

- Vos, N.J., Singh, N.A., Ross, D.A., Stavrinou, T.M., Orr, R., and Singh, M.A.F. (2005). Optimal Load for Increasing Muscle Power During Explosive Resistance Training in Older Adults. *The Journals of Gerontology*, 60(5), 638-647.
- Warburton, D.E.R., Nicol, C.W., and Bredin, S.S.D. (2006). Health benefits of physical activity: the evidence. *Canadian Medical Association Journal*, 174(6), 801-809.
- Wei, M., Kampert, J.B., Barlow C.E., Nichaman, M.Z., Gibbons, L.W., Paffenbarger, R.S., et al. (1999). Relationship between low cardiorespiratory fitness and mortality in normal-weight, overweight, and obese men. *The Journal of The American Medical Association*, 27, 1547-1553.
- Yeong, Y.S., Oh, S.W., Baik, H.W., Park, H.S., and Kim, W.Y. (2004). Alcohol consumption and the metabolic syndrome in Korean adults: the 1998 Korean National Health and Nutrition Examination Survey American. *Journal of Clinical Nutrition*, 80(1), 217-224.
- Zhang, J.Q., Ji, L.L., Fogt, D.L., and Fretwell, V.S. (2007). Effect of exercise duration on postprandial hypertriglyceridemia in men with metabolic syndrome. *Journal of Applied Physiology*, 103(4), 1339-1345.
- Zhou, B (2002). Predictive values of body mass index and waist circumference for risk factors of certain related diseases in Chinese adults: study on optimal cut-off points of body mass index and waist circumference in Chinese adults. Asia Pacific. *Journal of Clinical Nutrition*, 11(58), 5685-5692.
- Zoeller, R.F. (2007). Physical Activity: The Role of Physical Activity and Fitness in the Prevention and Management of Type 2 Diabetes Mellitus. *American Journal of Lifestyle Medicine*, 1(5), 344-350.