



REFERENCES

มหาวิทยาลัยพระนคร

REFERENCES

- [1] Bennett, J., & Brown, C. M. (2000). Use of herbal remedies by patients in a health maintenance organization. Journal of the American Pharmaceutical Association, 40 (3), 353-358.
- [2] Akerele, O. (1992). WHO guidelines for the assessment of herbal medicines. Fitoterapia, 63 (2), 99-104.
- [3] Kneedler, J. A., Sky, S. S., & Sexton, L. R. (1998). Understanding alpha-hydroxy acids. Dermatology Nursing, 10 (4), 247-266.
- [4] Bergfeld, W. F. (1999). A lifetime of healthy skin: Implications for women. International Journal of Fertility, 44 (2), 83-95.
- [5] Howard, P. C. et al. (2002). Alpha-hydroxy acids: Consideration of the biological effects and possible role in photocarcinogenesis. Journal of Food and Drug Analysis, 10 (4), 258-261.
- [6] Takahashi, M., & Machida, Y. (1985). The influence of hydroxy acids on the rheological properties of stratum corneum. Journal of the Society of Cosmetic Chemists, 36, 177-187.
- [7] Van-Scott, E. J., & Ruey, Y. J. (1984). Hyperkeratinization corneocyte cohesion and alpha-hydroxy acids. Journal of the American Academy of Dermatology, 11, 867-879.
- [8] Smith, W. P. (1996). Epidermal and dermal effects of topical lactic acid. Journal of the American Academy of Dermatology, 35, 388-391.
- [9] Berardesca, E., & Mailbach, H. (1995). AHA mechanisms of action. Cosmetics and Toiletries, 110, 30-31.
- [10] Smith, W. P. (1994). Hydroxy acids and skin aging. Cosmetics and Toiletries, 109, 41-48.
- [11] Fartasch, M., Teal, J., & Menon, G. K. (1997). Mode of action of glycolic acid on human stratum corneum: Ultrastructural and functional evaluation of the epidermal barrier. Archives of Dermatological Research, 289, 404-409.

- [12] Berardesca, E. et al (1997). Alpha hydroxyacids modulate stratum corneum barrier function. British Journal of Dermatology, 137, 934-938.
- [13] Lewis, Y. S., Neelakantan, S., & Bhatin, D. S. (1961). Determination of free and combined tartaric acid in plant products. Journal of Food Science (Mysore), 10, 49-50.
- [14] Lewis, Y. S., & Neelakantan, S. (1960). Nature of tartaric acid in tamarind. Journal of Food Science (Mysore), 9 (12), 405.
- [15] Hasan. K., Salam, A., & Wahid, M. A. (1978). A reinvestigation of tamarind fruit. Pakistan Journal of Science, 30 (6), 71-73.
- [16] Ghosh, G. P., & Pathak, A. K. (1981). Studies on complex formation of phosphoric acid with fruits, tartaric and citric acids. Proceedings of the National Academy of Sciences India Section A, 51 (2), 209-212.
- [17] Ranjan, S., Patnaik, K. K., & Laloraya, M. M. (1961). Enzymic conversion of meso-tartrate to dextro-tartrate in tamarind. Naturwissenschaften, 48, 406.
- [18] Verhaar, L. G. (1948). Tartaric acid and other constituents in the fruits of *Tamarindus indica*. Chronica Naturae, 104 (1), 8-11.
- [19] Sudborough, J. J., & Vridhachalam, P. N. (1920). Tartaric acid from tamarinds. Journal of Indian Institute of Science, 3, 61-80.
- [20] Marsden, F. (1923). Utilization of tamarind. Journal of Indian Institute of Science, 5, 157-162.
- [21] Institute of medplant, Mahidol University, Thailand. (no date). Tamarind plant. Retrieved on March 20, 2005 from: <http://www.medplant.mahidol.ac.th/pubhealth/%5Cpubhealth%5Ctamarin.html>
- [22] Bergfeld, W. et al. (1997). Improving the cosmetic appearance of photoaged skin with glycolic acid. Journal of the American Academy of Dermatology, 36, 1011-1013.
- [23] Vidt, D. G., & Bergfeld, W. F. (1997). Cosmetic use of alpha-hydroxy acids. Cleveland Clinic Journal of Medicine, 64, 327-329.

- [24] McCallion, R., & Li Wan Po, A. (1993). A dry and photo-aged skin: manifestations and management. International Journal of Clinical Pharmacology and Therapeutics, 18, 15-32.
- [25] Ditre, C. M. et al. (1996). Effects of α -hydroxy acids on photoaged skin: A pilot clinical, histologic, and ultrastructural study. Journal of the American Academy of Dermatology, 34, 187-195.
- [26] Bernstein, E. F. et al. (1997). Citric acid increases viable epidermal thickness and glycosaminoglycan content in sun-damaged skin. Dermatologic Surgery, 23, 689-694.
- [27] Donnelly, T. A. et al. (1997). Dermal matrix deposition in an in vitro skin model following administration of alpha hydroxy acid (AHA) formulations. The Journal of Investigative Dermatology, 104, 669.
- [28] Boisnic, S. et al. (1997). Long term culture of normal skin to test the efficacy of a hydroxy acid-containing cream. European Journal of Dermatology, 7, 271-273.
- [29] Morton, J. F. (March 7, 1999). Tamarind (tamarindus indica). Retrived on June 20, 2005 from: <http://www.hort.purdue.edu/newcrops/morton/tamarind.html>.
- [30] Viyoch, J., Klinthong, N., & Siripaisai, W. (2003). Development of oil-in water emulsion containing tamarind fruit pulp extract. Naresuan University Journal, 11 (3), 29-49.
- [31] Viyoch, J. et al. (2003). Formulation and development of a patch containing tamarind fruit extract by using the blended chitosan-starch as a rate-controlling matrix. International Journal of Cosmetic Science, 25, 113-125.
- [32] Abe, T. et al. (1980). Seasonal variations in skin temperature, skin pH, evaporative water loss and skin surface lipid values on human skin. Chemical and Pharmaceutical Bulletin, 28, 387-392.
- [33] Solomon, B. A., & Shalita, A. R. (1996). Effects of detergents on acne. Clinical Dermatol, 14 (1), 909-959.
- [34] CMP Medica Australia (September 16, 2003). Skin biology and structure. Retrieved on June 30, 2006 from: <http://www.mydr.com.au/default.asp?Article=3718>

- [35] Courage & Khazaka electronic GmbH (no date). Information and operating instructions for the Mexameter®MX 18. Germany: Courage & Khazaka electronic GmbH.
- [36] Courage & Khazaka electronic GmbH (no date). Information and operating instructions for the Skin-pH-Meter®PH 900 and Corneometer®CM 825. Germany: Courage & Khazaka electronic GmbH.
- [37] Courage & Khazaka electronic GmbH (no date). Information and operating instructions for the Cutometer®MPA 580. Germany: Courage & Khazaka electronic GmbH.
- [38] Courage & Khazaka electronic GmbH (no date). Information and operating instructions for the Tewameter®TM 300. Germany: Courage & Khazaka electronic GmbH.
- [39] Pinnagoda, J. et al, (1990). Guidelines for transepidermal water loss (TEWL) measurement. Contact Dermatitis, 22, 164-178.
- [40] Korting, H. C. et al, (1991). Changes in skin pH and resident flora by washing with synthetic detergent preparations at pH 5.5 and 8.5. Journal of the Society of Cosmetic Chemists, 42, 147-158.
- [41] Schreiner, V., Maerker, H., & Hoppe, U. (1996). Dependence of barrier repair in human skin on intra- and extracellular pH. The Journal of Investigative Dermatology, 106, 917.
- [42] Kawada, A. et al, (2002). A new approach to the evaluation of whitening effect of cosmetic using computer analysis of video-captured image. Journal of Dermatological Science, 29, 10-18.
- [43] Yilmaz, E., & Borchert, H. H. (2006). Effect of lipid-containing, positively charged nanoemulsions on skin hydration, elasticity and erythema-an in vivo study. International Journal of Pharmaceutics, 307, 232-238.
- [44] Fong, S. S. L., Hung, L. K., & Cheng, J. C. Y. (1997). The cutometer and ultrasonography in the assessment of postburn hypertrophic scar-a preliminary study. Burns, 23 (1), s12-s18.

- [45] Dobrev, H. P. (1999). In vivo study of skin mechanical properties in patients with systemic sclerosis. Journal of the American Academy of Dermatology, 40 (3), 436-342.
- [46] Draaijers, L. J. et al. (2004). Skin elasticity meter or subjective evaluation in scars: A reliability assessment. Burns, 30, 109-114.
- [47] Barel, A. O., & Clarys, P. (1995). Comparison of methods for measurement of transepidermal water loss. In: Serup, J. & Jemec, G. B. E. (Eds). Handbook of non-invasive methods and the skin. Boca Raton: CRC Press.
- [48] Elsner, P., Wilhelm, D., & Maibach, H. I. (1990). Mechanical properties of human forearm and vulvar skin. British Journal of Dermatology, 122, 607-614.
- [49] Yosipovitch, G. et al. (2000). Skin barrier properties in different body areas in neonates. Pediatrics, 106 (1), 105-108.
- [50] Sator, P. G., Schmidt, J. B., & Honigsmann, H. (2003). Comparison of epidermal hydration and skin surface lipids in healthy individuals and in patients with atopic dermatitis. Journal of the American Academy of Dermatology, 48 (3), 352-358.
- [51] Levin, J., & Maibach, H. (2005). The correlation between transepidermal water loss and percutaneous absorption: An overview. Journal of controlled release, 103, 291-299.
- [52] Gendler, E. C. (1997). Topical treatment of the aging face. Dermatologic Clinics, 15 (4), 561-567.
- [53] Greaves, M. W. (1990). Topical α -hydroxy acid derivatives for relieving dry itching skin. Cosmetics and Toiletries, 105, 61-64.
- [54] Andersen, F. A. (1998). Final report on the assessment of glycolic acid, ammonium, calcium, potassium, and sodium glycolate, methyl, ethyl, propyl, and butyl glycolates, and lactic acid, ammonium, calcium, potassium, sodium, and TEA lactates, methyl, ethyl, isopropyl, and butyl lactates, and lauryl, myristyl, and cetyl lactates. International Journal of Clinical Pharmacology, Therapy, and Toxicology, 17 (1), 1-241.

- [55] Van Scott, E. J. & Yu, R. J. (1974). Control of keratinization with a-hydroxy acids and related compounds. Archives of Dermatology, 110, 586-590.
- [56] Sexton, C. R., & Rubin, M. G. (1994). An overview of alpha-hydroxy acids. Dermatology Nursing, 6, 17-22.
- [57] Murad, H., Shamban, A. T., & Premo, P. S. (1995). The use of glycolic acid as a peeling agent. Dermatologic Clinics, 13, 285-307.
- [58] Lavker, R. M., Kaidbey, K., & Leyden, J. J. (1992). Effects of topical ammonium lactate on cutaneous atrophy from a potent topical corticosteroid. Journal of the American Academy of Dermatology, 26, 535-544.
- [59] Thibault, P. K., Wlodarczyk, J. & Wenck, A. A. (1998). Double-blind randomized clinical trial on the effectiveness of a daily glycolic acid 5% formulation in the treatment of photoaging. Dermatologic Surgery, 24, 573-578.
- [60] Elson, M. L. (1992). The utilization of glycolic acid in photoaging. Cosmetic Dermatology, 5, 36.
- [61] Newman, N. et al. (1996). Clinical improvement of photoaged skin with 50% glycolic acid. Dermatologic Surgery, 22, 455-460.
- [62] Van Scott, E. J., Ditre, C. M., & Yu, R. J. (1996). Alpha-hydroxy acids in the treatment of signs of photoaging. Clinics in Dermatology, 14, 217-226.
- [63] Van Scott, E. J., & Yu, R. J. (1989). Alpha-hydroxy acids: Therapeutic potentials. Canada Journal Dermatology, 1 (5), 108-112.
- [64] Sexton, C. R., & Rubin, M. G. (1994). An overview of alpha-hydroxy acids. Dermatology Nursing, 6, 17-22.
- [65] Moy, L. S., Murad, H., & Moy, R. J. (1993). Glycolic acid peels for the treatment of wrinkles and photoaging. Journal of Dermatologic Surgery, 19, 243.
- [66] Kurtzweil, P. (May, 1999). Alpha hydroxy acids for skin care. Retrieved on May 15, 2005 from: <http://vm.cfsan.fda.gov/~dms/fdacaha.html>.
- [67] Friedman, L. M., Furberg, C. D., & Demets, D. L. (1998). Fundamental of clinical trials (3rd ed). Springer: Mosby.

- [68] Margarone, J. E. et al. (1995). A comparison of crossover versus parallel-group design in the evaluation of analgesic efficacy after molar extraction. Clinical pharmacology and therapeutics, 58 (4), 453-458.
- [69] Richens, A. (2001). Proof of efficacy trials: Cross-over versus parallel-group. Epilepsy Research, 45, 434-437.
- [70] Byar D. P. et al. (1976). Randomized clinical trials: Perspectives on some recent ideas. The New England Journal of Medicine, 295, 74-80.
- [71] Moher, D., Schulz, K. F., & Altman, D. G. (2001). The CONSORT statement: Revised recommendations for improving the quality of reports of parallel group randomized trials. Annals of Internal Medicine, 134, 657-662.
- [72] Keech, A. (1998). Random allocation, introduction to clinical trials, clinical trials research methodology, and statistical methods in clinical trials. The ICH GCP guidelines. Hong Kong: The Clinical Trials Centre.
- [73] Pocock, S. (1984). Clinical trials: A practical approach. New York: John Wiley.
- [74] Leyden, J. J., & Rawlings, A. V. (Eds) (2002). Skin moisturizer : In cosmetic science and technology series. New York: Marcel Dekker.
- [75] Schulz, K. F. et al. (1995). Empirical evidence of bias. Dimensions of methodological quality associated with estimates of treatment effects in controlled trials. Journal of the American Medical Association, 273, 408-412.
- [76] Rosner B. (1995). Fundamentals of biostatistics (4th ed.). USA: Duxbury Press.
- [77] Greenland, S., & Finkle, W. D. (1995). A critical look at methods for handling missing covariates in epidemiologic regression analysis. American Journal of Epidemiology, 142, 1225-1264.
- [78] Edison, B. L. et al. (2004). A polyhydroxy acid skin care regimen provides antiaging effects comparable to an alpha-hydroxyacid regimen. Cutis, 73 (2), 14-17.
- [79] Kaidbey, K. et al. (2003). Topical glycolic acid enhances photodamage by ultraviolet light. Photodermatology, Photoimmunology & Photomedicine, 19 (1), 21-27.

- [80] Stiller, M. J. et al. (1996). Topical 8% glycolic acid and 8% L-lactic acid creams for the treatment of photodamaged skin: a double-blind vehicle-controlled clinical trial. Archives of Dermatology, 132 (6), 631-636.
- [81] Kempers, S. (1998). An evaluation of the effect of an alpha hydroxy acid-blend skin cream in the cosmetic improvement of symptoms of moderate to severe xerosis, epidermolytic hyperkeratosis, and ichthyosis. Cutis, 61 (6), 347-350.
- [82] Haddad, A. L. et al. (2003). A clinical, prospective, randomized, double-blind trial comparing skin whitening complex with hydroquinone vs. placebo in the treatment of melasma. International Journal of Dermatology, 42 (2), 153-156.
- [83] Kawada, A. et al. (2002). A new approach to the evaluation of whitening effect of a cosmetic using computer analysis of video-captured image. Journal of Dermatological Science, 29, 10-18.
- [84] Briganti, S., Camera, E., & Picardo, M. (2003). Chemical and instrumental approaches to treat hyperpigmentation. Pigment Cell Research, 16, 101-109.
- [85] Shu, Y. T. et al. (2005). The investigation on the skin whitening efficacy of Chinese herb extracts using the colour physics technology. Journal of Dermatological Science, 37 (3), 177-179.
- [86] Tsai, T. F. et al. (2000). Effect of glycolic acid on light-induced skin pigmentation in Asians and Caucasian subjects. Journal of the American Academy of Dermatology, 43 (2), 238-243.
- [87] Takiwaki, H., Overgaard, L., & Serup, J. (1994). Comparison of narrow-band reflectance spectrophotometric and tristimulus colorimetric measurements of skin color. Twenty-three anatomical sites evaluated by the Deraspectrometer and the Chroma Meter CR-200. Skin Pharmacology: the Official Journal of the Skin Pharmacology Society, 7 (4), 217-225.
- [88] Clarys, P. et al. (2000). Skin color measurements: comparison between three instruments: The Chromameter (R), the DermaSpectrometer (R) and the Mexameter (R). Skin Research and Technology, 6 (4), 230-238.

- [89] Shriver, M. D., & Parra, E. J. (2000). Comparison of narrow-band reflectance spectroscopy and tristimulus colorimetry for measurements of skin and hair color in persons of different biological ancestry. American Journal of Physical Anthropology, 112 (1), 17-27.
- [90] Yoshimura, K. et al. (2001). Usefulness of a narrow-band reflectance spectrophotometer in evaluating effects of depigmenting treatment. Aesthetic Plastic Surgery, 25 (2), 129-133.

