



BIBLIOGRAPHY

- [1] Bureau of policy and strategy ministry of public health, Thailand. (March 18, 2005). Public health statistics 2003. Retrieved on March 22, 2005. from: <http://203.157.19.191/index-stat46.htm>
- [2] Murdoch, D., & Heel, R. C. (1991). Amlodipine a review of its pharmacodynamic and pharmacokinetic properties, and therapeutic use in cardiovascular disease. Drugs, 41 (3), 478-505.
- [3] Budavari, S. et al. (Ed.) (1996). The merck index, an encyclopedia of chemicals, drugs, and biologicals (12th ed.). USA: Merck & Co.
- [4] Haria, M., & Wagstaff, A. J. (1995). Amlodipine a reappraisal of its pharmacological properties and therapeutic use in cardiovascular disease. Drugs, 50 (3), 560-586.
- [5] Pfizer Labs. (2000). Norvasc[®] (amlodipine besylate) Tablets. USA: Pfizer.
- [6] Meredith, P. A., & Elliott, H. L. (1992). Clinical pharmacokinetics of amlodipine. Clin Pharmacokinet, 22 (1), 22-31.
- [7] Faulkner, J. K., Hayden, M. L., & Taylor, T. (1989). Absorption of amlodipine unaffected by food. Solid dose equivalent to solution dose. Arzeim-Forsch./Drug Res, 39 (2), 799-801.
- [8] Monkman, S. C. et al. (1996). Automated gas chromatographic assay for amlodipine in plasma and gingival crevicular fluid. J Chromatogr B, 678, 360-364.
- [9] Faulkner, J. K. et al. (1986). The pharmacokinetics of amlodipine in Healthy volunteers after single intravenous and oral doses and after 14 repeated oral doses given once daily. Br J Clin Pharmacol, 22 (1), 21-25.
- [10] Tatar, S., & Atmaca, S. (2001). Determination of amlodipine in human plasma by high-performance liquid chromatography with fluorescence detection. J Chromatogr B, 758, 305-310.

- [11] Argekar, A. P., & Power, S. G. (2000). Simultaneous determination of atenolol and amlodipine in tablet by high-performance thin-layer chromatography. J Pharm Biomed Anal, 21, 1137-1142.
- [12] Schmuff, N. R., & Lunn, G. (Ed.) (1996). HPLC methods for pharmaceutical analysis. USA: John Wiley & Sons.
- [13] Patel, Y. P. et al. (1998). Isocratic, simultaneous reversed-phase high-performance liquid chromatographic estimation of six drugs for combined hypertension therapy. J Chromatogr A, 828, 283-286.
- [14] Josefsson, M., Zackrisson, A. L., & Norlander, B. (1995). Sensitive high-performance liquid chromatographic analysis of amlodipine in human plasma with amperometric detection and a single-step solid-phase sample preparation. J Chromatogr B, 672, 310-313.
- [15] Shimooka, K., Sawada, Y., & Tatamatsu, H. (1989). Analysis of amlodipine in serum by a sensitive high-performance liquid chromatographic method with amperometric detection. J Pharm Biomed Anal, 7 (11), 1267-1272.
- [16] Carvalho, M. et al. (2001). Amlodipine bioequivalence study: Quantification by liquid chromatography coupled to tandem mass spectrometry. Biopharm Drug Dispos, 22, 383-390.
- [17] Zhong, D. et al. (2001). Applications of liquid chromatography-tandem mass spectrometry in drug and biomedical analyses. Clin Chim Acta, 313, 147-150.
- [18] Streeb, B. et al. (2002). Enantiomeric determination of amlodipine in human plasma by liquid chromatography coupled to tandem mass spectrometry. J Biochem Biophys Methods, 54, 357-368.
- [19] Drug Control Division Food and Drug administration. (2004). Thailand guidelines for the conduct of bioavailability and bioequivalence studies. Nonthaburi: Drug control division Food and Drug administration, Ministry of Public Health.
- [20] U.S. Department of Health and Human Services. (2001). Guidance for industrial Bioanalytical method validation. USA: U.S. Food and Drug Administration.

- [21] Webster, J. et al. (1987). Once daily amlodipine in the treatment of mild to moderate hypertension. Br J Clin Pharmacol, 24, 713-719.
- [22] Bainbridge, A. D. (1993). A comparative assessment of amlodipine and felodipine ER: pharmacokinetic and pharmacodynamic indices. Eur J Clin Pharmacol, 45, 425-430.
- [23] Josefsson, M., & Norlander, B. (1996). Coupled-column chromatography on a Chiral-AGP phase for determination of amlodipine enantiomers in human plasma: An HPLC assay with electrochemical detection. J Pharm Biomed Anal, 15, 267-277.
- [24] Kurtz, G. S. et al. (1999). Limited-sampling strategy models for estimating the area under the plasma concentration-time curve for amlodipine. Eur J Clin Pharmacol, 55, 651-657.
- [25] Aktas, E. S., Ersoy, L., & Sagirli, O. (2003). A new spectrofluorimetric method for the determination of lisinopril in tablets. Il Farmaco, 58, 165-168.