



บรรณานุกรม

มหาวิทยาลัยราชภัฏบรจรม

บรรณานุกรม

- เมธี วงศ์ศิริสุวรรณ. (2551). เมื่อสภาวะแวดล้อมและยีนมาบรรจบกัน: สาเหตุของการเกิดโรคพาร์กินสัน. *Medical progress CME*, 7(3), 34-39.
- Abeliovich, A., Schmitz, Y., Farinas, I., Choi-Lundberg, D., Ho, W. H., Castillo, P. E., et al. (2000). Mice lacking alpha-synuclein display functional deficits in the nigrostriatal dopamine system. *Neuron*, 25(1), 239-252.
- Aman, M. J., Tayebi, N., Obiri, N. I., Puri, R. K., Modi, W. S. and Leonard, W. J. (1996). cDNA cloning and characterization of the human interleukin 13 receptor alpha chain. *J Biol Chem*, 271(46), 29265-29270.
- Andersen, K., Launer, L. J., Dewey, M. E., Letenneur, L., Ott, A., Copeland, J. R., et al. (1999). Gender differences in the incidence of AD and vascular dementia: The EURODEM Studies. EURODEM Incidence Research Group. *Neurology*, 53(9), 1992-1997.
- Ariga, T., Kiso, M., Hasegawa, A. and Miyatake, T. (2001). Gangliosides inhibit the release of interleukin-1beta in amyloid beta-protein-treated human monocytic cells. *J Mol Neurosci*, 17(3), 371-377.
- Bellani, S., Sousa, V. L., Ronzitti, G., Valtorta, F., Meldolesi, J. and Chieregatti, E. (2010). The regulation of synaptic function by alpha-synuclein. *Commun Integr Biol*, 3(2), 106-109.
- Bennett, D. A., Schneider, J. A., Tang, Y., Arnold, S. E. and Wilson, R. S. (2006). The effect of social networks on the relation between Alzheimer's disease pathology and level of cognitive function in old people: a longitudinal cohort study. *Lancet Neurol*, 5(5), 406-412.
- Bermejo-Pareja, F., Benito-Leon, J., Vega, S., Medrano, M. J. and Roman, G. C. (2008). Incidence and subtypes of dementia in three elderly populations of central Spain. *J Neurol Sci*, 264(1-2), 63-72.
- Bialystok, E., Craik, F. I. M. and Freedman, M. (2007). Bilingualism as a protection against the onset of symptoms of dementia. *Neuropsychologia*, 45(2), 459-464.

- Block, M. L. and Hong, J.-S. (2005). Microglia and inflammation-mediated neurodegeneration: Multiple triggers with a common mechanism. *Progress in Neurobiology*, 76(2), 77-98.
- Bonin-Guillaume, S., Zekry, D., Giacobini, E., Gold, G. and Michel, J. P. (2005). The economical impact of dementia. *Presse Med*, 34(1), 35-41.
- Brookmeyer, R., Gray, S. and Kawas, C. (1998). Projections of Alzheimer's disease in the United States and the public health impact of delaying disease onset. *Am J Public Health*, 88(9), 1337-1342.
- Brookmeyer, R., Johnson, E., Ziegler-Graham, K. and Arrighi, H. M. (2007). Forecasting the global burden of Alzheimer's disease. *Alzheimer's and Dementia*, 3(3), 186-191.
- Cabin, D. E., Shimazu, K., Murphy, D., Cole, N. B., Gottschalk, W., McIlwain, K. L., et al. (2002). Synaptic vesicle depletion correlates with attenuated synaptic responses to prolonged repetitive stimulation in mice lacking alpha-synuclein. *J Neurosci*, 22(20), 8797-8807.
- Cantu, D., Fulton, R. E., Drechsel, D. A. and Patel, M. (2011). Mitochondrial aconitase knockdown attenuates paraquat-induced dopaminergic cell death via decreased cellular metabolism and release of iron and H₂O₂. *J Neurochem*, 118(1), 79-92.
- Caput, D., Laurent, P., Kaghad, M., Lelias, J. M., Lefort, S., Vita, N., et al. (1996). Cloning and characterization of a specific interleukin (IL) -13 binding protein structurally related to the IL-5 receptor alpha chain. *J Biol Chem*, 271(28), 16921-16926.
- Chang, P.-A., Wu, Y.-J., Li, W. and Leng, X.-F. (2006). Effect of carbamate esters on neurite outgrowth in differentiating human SK-N-SH neuroblastoma cells. *Chemico-Biological Interactions*, 159(1), 65-72.
- Chaouchi, N., Wallon, C., Goujard, C., Tertian, G., Rudent, A., Caput, D., et al. (1996). Interleukin-13 inhibits interleukin-2-induced proliferation and protects chronic lymphocytic leukemia B cells from in vitro apoptosis. *Blood*, 87(3), 1022-1029.
- Chen, L., Yoo, S.-E., Na, R., Liu, Y. and Ran, Q. (2012). Cognitive impairment and increased A β levels induced by paraquat exposure are attenuated by enhanced removal of mitochondrial H₂O₂. *Neurobiology of Aging*, 33(2), 432.e415-432.e426.

- Chiou, W. F. (2006). Effect of Abeta exposure on the mRNA expression patterns of voltage-sensitive calcium channel alpha 1 subunits (alpha 1A-alpha 1D) in human SK-N-SH neuroblastoma. *Neurochem Int*, 49(3), 256-261.
- Cho, W., Kim, Y., Jeoung, D. I., Kim, Y. M. and Choe, J. (2011). IL-4 and IL-13 suppress prostaglandins production in human follicular dendritic cells by repressing COX-2 and mPGES-1 expression through JAK1 and STAT6. *Mol Immunol*, 48(6-7), 966-972.
- Choi, W. S., Abel, G., Klintworth, H., Flavell, R. A. and Xia, Z. (2010). JNK3 mediates paraquat- and rotenone-induced dopaminergic neuron death. *J Neuropathol Exp Neurol*, 69(5), 511-520.
- Chun, W. and Johnson, G. V. (2007). The role of tau phosphorylation and cleavage in neuronal cell death. *Front Biosci*, 12, 733-756.
- Cookson, M. R. (2009). alpha-Synuclein and neuronal cell death. *Mol Neurodegener*, 4, 9.
- Cottrez, F., Hurst, S. D., Coffman, R. L. and Groux, H. (2000). T regulatory cells 1 inhibit a Th2-specific response in vivo. *J Immunol*, 165(9), 4848-4853.
- Cribbs, D. H. (2010). Abeta DNA Vaccination for Alzheimer's Disease: Focus on Disease Prevention. *CNS Neurol Disord Drug Targets*, 9(2), 207-216.
- da Costa, C. A., Ancolio, K. and Checler, F. (2000). Wild-type but not Parkinson's disease-related ala-53 --> Thr mutant alpha -synuclein protects neuronal cells from apoptotic stimuli. *J Biol Chem*, 275(31), 24065-24069.
- Davis, E., Krishnan, K., Yan, H., Newcomb, E. W. and Krolewski, J. J. (1996). A mutant form of p135tyk2, an interferon-alpha inducible tyrosine kinase, suppresses the transformed phenotype of Daudi cells. *Leukemia*, 10(3), 543-551.
- Dawson, T. M. and Dawson, V. L. (2003a). Molecular pathways of neurodegeneration in Parkinson's disease. *Science*, 302(5646), 819-822.
- Dawson, T. M. and Dawson, V. L. (2003b). Rare genetic mutations shed light on the pathogenesis of Parkinson disease. *J Clin Invest*, 111(2), 145-151.
- de Lau, L. M. and Breteler, M. M. (2006). Epidemiology of Parkinson's disease. *Lancet Neurol*, 5(6), 525-535.

- de Waal Malefyt, R., Abrams, J. S., Zurawski, S. M., Lecron, J. C., Mohan-Peterson, S., Sanjanwala, B., et al. (1995). Differential regulation of IL-13 and IL-4 production by human CD8+ and CD4+ Th0, Th1 and Th2 T cell clones and EBV-transformed B cells. *Int Immunol*, 7(9), 1405-1416.
- de Waal Malefyt, R., Figdor, C. G., Huijbens, R., Mohan-Peterson, S., Bennett, B., Culpepper, J., et al. (1993). Effects of IL-13 on phenotype, cytokine production, and cytotoxic function of human monocytes. Comparison with IL-4 and modulation by IFN-gamma or IL-10. *J Immunol*, 151(11), 6370-6381.
- Di Carlo, A., Baldereschi, M., Amaducci, L., Lepore, V., Bracco, L., Maggi, S., et al. (2002). Incidence of dementia, Alzheimer's disease, and vascular dementia in Italy. The ILSA Study. *J Am Geriatr Soc*, 50(1), 41-48.
- Diaz-Cazorla, M., Perez-Sala, D., Ros, J., Jimenez, W., Fresno, M. and Lamas, S. (1999). Regulation of cyclooxygenase-2 expression in human mesangial cells--transcriptional inhibition by IL-13. *Eur J Biochem*, 260(1), 268-274.
- Doucet, C., Brouty-Boye, D., Pottin-Clemenceau, C., Canonica, G. W., Jasmin, C. and Azzarone, B. (1998). Interleukin (IL) 4 and IL-13 act on human lung fibroblasts. Implication in asthma. *J Clin Invest*, 101(10), 2129-2139.
- Dunnett, S. B. and Bjorklund, A. (1999). Prospects for new restorative and neuroprotective treatments in Parkinson's disease. *Nature*, 399(6738 Suppl), A32-39.
- Farlow, M. R. and Brosch, J. R. (2013). Immunotherapy for Alzheimer's Disease. *Neurologic Clinics*, 31(3), 869-878.
- Forloni, G., Bertani, I., Calella, A. M., Thaler, F. and Invernizzi, R. (2000). Alpha-synuclein and Parkinson's disease: selective neurodegenerative effect of alpha-synuclein fragment on dopaminergic neurons in vitro and in vivo. *Ann Neurol*, 47(5), 632-640.
- Fortin, D. L., Troyer, M. D., Nakamura, K., Kubo, S., Anthony, M. D. and Edwards, R. H. (2004). Lipid rafts mediate the synaptic localization of alpha-synuclein. *J Neurosci*, 24(30), 6715-6723.

- Franco, R., Li, S., Rodriguez-Rocha, H., Burns, M. and Panayiotidis, M. I. (2010). Molecular mechanisms of pesticide-induced neurotoxicity: Relevance to Parkinson's disease. *Chem Biol Interact*, 188(2), 289-300.
- Fu, X. Y., Kessler, D. S., Veals, S. A., Levy, D. E. and Darnell, J. E., Jr. (1990). ISGF3, the transcriptional activator induced by interferon alpha, consists of multiple interacting polypeptide chains. *Proc Natl Acad Sci U S A*, 87(21), 8555-8559.
- Gabbita, S. P., Butterfield, D. A., Hensley, K., Shaw, W. and Carney, J. M. (1997). Aging and caloric restriction affect mitochondrial respiration and lipid membrane status: an electron paramagnetic resonance investigation. *Free Radic Biol Med*, 23(2), 191-201.
- Garcia-Alloza, M., Borrelli, L. A., Rozkalne, A., Hyman, B. T. and Bacskai, B. J. (2007). Curcumin labels amyloid pathology in vivo, disrupts existing plaques, and partially restores distorted neurites in an Alzheimer mouse model. *J Neurochem*, 102(4), 1095-1104.
- Garwood, E. R., Bekele, W., McCulloch, C. E. and Christine, C. W. (2006). Amphetamine exposure is elevated in Parkinson's disease. *NeuroToxicology*, 27(6), 1003-1006.
- Gelb, D. J., Oliver, E. and Gilman, S. (1999). Diagnostic criteria for Parkinson disease. *Archives of neurology*, 56(1), 33-39.
- Ghochikyan, A., Mkrtychyan, M., Petrushina, I., Movsesyan, N., Karapetyan, A., Cribbs, D. H., et al. (2006). Prototype Alzheimer's disease epitope vaccine induced strong Th2-type anti-Abeta antibody response with Alum to Quil A adjuvant switch. *Vaccine*, 24(13), 2275-2282.
- Giasson, B. I., Ischiropoulos, H., Lee, V. M. and Trojanowski, J. Q. (2002). The relationship between oxidative/nitrative stress and pathological inclusions in Alzheimer's and Parkinson's diseases. *Free Radic Biol Med*, 32(12), 1264-1275.
- Gray, J. P., Heck, D. E., Mishin, V., Smith, P. J., Hong, J. Y., Thiruchelvam, M., et al. (2007). Paraquat increases cyanide-insensitive respiration in murine lung epithelial cells by activating an NAD (P) H:paraquat oxidoreductase: identification of the enzyme as thioredoxin reductase. *J Biol Chem*, 282(11), 7939-7949.

- Gray, S. L., Anderson, M. L., Crane, P. K., Breitner, J. C., McCormick, W., Bowen, J. D., et al. (2008). Antioxidant vitamin supplement use and risk of dementia or Alzheimer's disease in older adults. *J Am Geriatr Soc*, 56(2), 291-295.
- Groner, B., Altioek, S. and Meier, V. (1994). Hormonal regulation of transcription factor activity in mammary epithelial cells. *Mol Cell Endocrinol*, 100(1-2), 109-114.
- Gurunathan, S., Wu, C. Y., Freidag, B. L. and Seder, R. A. (2000). DNA vaccines: a key for inducing long-term cellular immunity. *Curr Opin Immunol*, 12(4), 442-447.
- Hayden, K. M., Norton, M. C., Darcey, D., Ostbye, T., Zandi, P. P., Breitner, J. C., et al. (2010). Occupational exposure to pesticides increases the risk of incident AD: the Cache County study. *Neurology*, 74(19), 1524-1530.
- Hebert, L. E., Scherr, P. A., Bienias, J. L., Bennett, D. A. and Evans, D. A. (2003). Alzheimer disease in the US population: prevalence estimates using the 2000 census. *Arch Neurol*, 60(8), 1119-1122.
- Hirsch, E. C. and Hunot, S. (2009). Neuroinflammation in Parkinson's disease: a target for neuroprotection? *Lancet Neurol*, 8(4), 382-397.
- Hoehn, M. M. and Yahr, M. D. (2001). Parkinsonism: onset, progression, and mortality. 1967. *Neurology*, 57(10 Suppl 3), S11-26.
- Humphrey, M. L., Cole, M. P., Pendergrass, J. C. and Kinningham, K. K. (2005). Mitochondrial Mediated Thimerosal-Induced Apoptosis in a Human Neuroblastoma Cell Line (SK-N-SH). *NeuroToxicology*, 26(3), 407-416.
- Hunot, S., Vila, M., Teismann, P., Davis, R. J., Hirsch, E. C., Przedborski, S., et al. (2004). JNK-mediated induction of cyclooxygenase 2 is required for neurodegeneration in a mouse model of Parkinson's disease. *Proc Natl Acad Sci U S A*, 101(2), 665-670.
- Iqbal, K., Alonso Adel, C., Chen, S., Chohan, M. O., El-Akkad, E., Gong, C. X., et al. (2005). Tau pathology in Alzheimer disease and other tauopathies. *Biochim Biophys Acta*, 1739(2-3), 198-210.

- Iwatsubo, T., Odaka, A., Suzuki, N., Mizusawa, H., Nukina, N. and Ihara, Y. (1994).
Visualization of A beta 42 (43) and A beta 40 in senile plaques with end-specific A
beta monoclonals: evidence that an initially deposited species is A beta 42 (43).
Neuron, 13(1), 45-53.
- Izuhara, K., Arima, K., Kanaji, S., Kanaji, T. and Ohta, S. (2007). IL-13 and its Signal Pathway:
Promising Targets in the Development of a Therapeutic Agent for Bronchial
Asthma. *Current Signal Transduction Therapy*, 2(1), 31-40.
- Jo, E., McLaurin, J., Yip, C. M., St George-Hyslop, P. and Fraser, P. E. (2000). alpha-
Synuclein membrane interactions and lipid specificity. *J Biol Chem*, 275(44),
34328-34334.
- Johnston, J. A., Ward, C. L. and Kopito, R. R. (1998). Aggresomes: a cellular response to
misfolded proteins. *J Cell Biol*, 143(7), 1883-1898.
- Kalaria, R. N. (1999). Microglia and Alzheimer's disease. *Curr Opin Hematol*, 6(1), 15-24.
- Kawahara, K., Suenobu, M., Yoshida, A., Koga, K., Hyodo, A., Ohtsuka, H., et al. (2012).
Intracerebral microinjection of interleukin-4/interleukin-13 reduces β -amyloid
accumulation in the ipsilateral side and improves cognitive deficits in young
amyloid precursor protein 23 mice. *Neuroscience*, 207, 243-260.
- Kawas, C. H. (2006). Medications and diet: protective factors for AD? *Alzheimer Dis Assoc
Disord*, 20(3 Suppl 2), S89-96.
- Kim, D. W., Uetsuki, T., Kaziro, Y., Yamaguchi, N. and Sugano, S. (1990). Use of the human
elongation factor 1 alpha promoter as a versatile and efficient expression system.
Gene, 91(2), 217-223.
- Kim, E.-H., Lee, M.-J., Kim, I.-H., Pyo, S.-N., Choi, K.-T. and Rhee, D.-K. (2010). Anti-
apoptotic Effects of Red Ginseng on Oxidative Stress Induced by Hydrogen
Peroxide in SK-N-SH Cells. *Journal of Ginseng Research*, 34(2), 138.
- Kim, H. D., Cao, Y., Kong, F. K., Van Kampen, K. R., Lewis, T. L., Ma, Z., et al. (2005).
Induction of a Th2 immune response by co-administration of recombinant
adenovirus vectors encoding amyloid beta-protein and GM-CSF. *Vaccine*, 23(23),
2977-2986.

- Kimura, M., Takatsuki, A. and Yamaguchi, I. (1994). Blastocidin S deaminase gene from *Aspergillus terreus* (BSD) : a new drug resistance gene for transfection of mammalian cells. *Biochimica et Biophysica Acta (BBA) - Gene Structure and Expression*, 1219(3), 653-659.
- Klegeris, A. and McGeer, P. L. (1997). beta-amyloid protein enhances macrophage production of oxygen free radicals and glutamate. *J Neurosci Res*, 49(2), 229-235.
- Kutzler, M. A., Robinson, T. M., Chattergoon, M. A., Choo, D. K., Choo, A. Y., Choe, P. Y., et al. (2005). Coimmunization with an optimized IL-15 plasmid results in enhanced function and longevity of CD8 T cells that are partially independent of CD4 T cell help. *J Immunol*, 175(1), 112-123.
- LaFerla, F. M., Green, K. N. and Oddo, S. (2007). Intracellular amyloid-beta in Alzheimer's disease. *Nat Rev Neurosci*, 8(7), 499-509.
- Lang, A. E. and Lozano, A. M. (1998). Parkinson's disease. First of two parts. *N Engl J Med*, 339(15), 1044-1053.
- Ling, Y., Morgan, K. and Kalsheker, N. (2003). Amyloid precursor protein (APP) and the biology of proteolytic processing: relevance to Alzheimer's disease. *Int J Biochem Cell Biol*, 35(11), 1505-1535.
- Litteljohn, D., Mangano, E., Clarke, M., Bobyn, J., Moloney, K. and Hayley, S. (2010). Inflammatory mechanisms of neurodegeneration in toxin-based models of Parkinson's disease. *Parkinsons Dis*, 2011, 713517.
- Liu, M. A. (2011). DNA vaccines: an historical perspective and view to the future. *Immunol Rev*, 239(1), 62-84.
- Manna, S. K. and Aggarwal, B. B. (1998). IL-13 suppresses TNF-induced activation of nuclear factor-kappa B, activation protein-1, and apoptosis. *J Immunol*, 161(6), 2863-2872.
- Manning-Bog, A. B., McCormack, A. L., Li, J., Uversky, V. N., Fink, A. L. and Di Monte, D. A. (2002). The herbicide paraquat causes up-regulation and aggregation of alpha-synuclein in mice: paraquat and alpha-synuclein. *J Biol Chem*, 277(3), 1641-1644.

- Manning-Bog, A. B., McCormack, A. L., Purisai, M. G., Bolin, L. M. and Di Monte, D. A. (2003). Alpha-synuclein overexpression protects against paraquat-induced neurodegeneration. *J Neurosci*, 23(8), 3095-3099.
- Martin, F. L., Williamson, S. J., Paleologou, K. E., Allsop, D. and El-Agnaf, O. M. (2004). Alpha-synuclein and the pathogenesis of Parkinson's disease. *Protein Pept Lett*, 11(3), 229-237.
- Masliah, E., Rockenstein, E., Veinbergs, I., Mallory, M., Hashimoto, M., Takeda, A., et al. (2000). Dopaminergic loss and inclusion body formation in alpha-synuclein mice: implications for neurodegenerative disorders. *Science*, 287(5456), 1265-1269.
- Mateen, I. and Irshad, S. (2011). A Review on DNA Vaccines. *Journal of Health Science*, 1(1), 1-7.
- Mathews, P. M. and Nixon, R. A. (2003). Setback for an Alzheimer's disease vaccine: lessons learned. *Neurology*, 61(1), 7-8.
- McKenzie, A. N., Culpepper, J. A., de Waal Malefyt, R., Briere, F., Punnonen, J., Aversa, G., et al. (1993). Interleukin 13, a T-cell-derived cytokine that regulates human monocyte and B-cell function. *Proc Natl Acad Sci U S A*, 90(8), 3735-3739.
- McNaught, K. S., Shashidharan, P., Perl, D. P., Jenner, P. and Olanow, C. W. (2002). Aggresome-related biogenesis of Lewy bodies. *Eur J Neurosci*, 16(11), 2136-2148.
- Meek, P. D., McKeithan, K. and Schumock, G. T. (1998). Economic considerations in Alzheimer's disease. *Pharmacotherapy*, 18(2 Pt 2), 68-73; discussion 79-82.
- Miller, D. W., Hague, S. M., Clarimon, J., Baptista, M., Gwinn-Hardy, K., Cookson, M. R., et al. (2004). Alpha-synuclein in blood and brain from familial Parkinson disease with SNCA locus triplication. *Neurology*, 62(10), 1835-1838.
- Miloux, B., Laurent, P., Bonnin, O., Lupker, J., Caput, D., Vita, N., et al. (1997). Cloning of the human IL-13 R alpha1 chain and reconstitution with the IL4R alpha of a functional IL-4/IL-13 receptor complex. *FEBS Lett*, 401(2-3), 163-166.
- Minty, A., Ferrara, P. and Caput, D. (1997). Interleukin-13 effects on activated monocytes lead to novel cytokine secretion profiles intermediate between those induced by interleukin-10 and by interferon-gamma. *Eur Cytokine Netw*, 8(2), 189-201.

- Molsa, P. K., Marttila, R. J. and Rinne, U. K. (1995). Long-term survival and predictors of mortality in Alzheimer's disease and multi-infarct dementia. *Acta Neurol Scand*, 91(3), 159-164.
- Moongkarndi, P., Srisawat, C., Saetun, P., Jantaravinid, J., Peerapittayamongkol, C., Soi-ampornkul, R., et al. (2010). Protective effect of mangosteen extract against beta-amyloid-induced cytotoxicity, oxidative stress and altered proteome in SK-N-SH cells. *J Proteome Res*, 9(5), 2076-2086.
- Moore, A. H. and O'Banion, M. K. (2002). Neuroinflammation and anti-inflammatory therapy for Alzheimer's disease. *Advanced Drug Delivery Reviews*, 54(12), 1627-1656.
- Morris, M. C., Evans, D. A., Bienias, J. L., Scherr, P. A., Tangney, C. C., Hebert, L. E., et al. (2004). Dietary niacin and the risk of incident Alzheimer's disease and of cognitive decline. *J Neurol Neurosurg Psychiatry*, 75(8), 1093-1099.
- Mudher, A. and Lovestone, S. (2002). Alzheimer's disease-do taoists and baptists finally shake hands? *Trends Neurosci*, 25(1), 22-26.
- Nistor, M., Don, M., Parekh, M., Sarsoza, F., Goodus, M., Lopez, G. E., et al. (2007). Alpha- and beta-secretase activity as a function of age and beta-amyloid in Down syndrome and normal brain. *Neurobiol Aging*, 28(10), 1493-1506.
- Nostrandt, A. C. and Ehrich, M. (1992). Development of a model cell culture system in which to study early effects of neuropathy-inducing organophosphorus esters. *Toxicology Letters*, 60(1), 107-114.
- O'Shea, J. J. (1997). Jaks, STATs, cytokine signal transduction, and immunoregulation: are we there yet? *Immunity*, 7(1), 1-11.
- Obiri, N. I., Murata, T., Debinski, W. and Puri, R. K. (1997). Modulation of interleukin (IL) -13 binding and signaling by the gammac chain of the IL-2 receptor. *J Biol Chem*, 272(32), 20251-20258.
- Oda, T., Kosuge, Y., Arakawa, M., Ishige, K. and Ito, Y. (2008). Distinct mechanism of cell death is responsible for tunicamycin-induced ER stress in SK-N-SH and SH-SY5Y cells. *Neurosci Res*, 60(1), 29-39.

- Ogata, H., Ford, D., Kouttab, N., King, T. C., Vita, N., Minty, A., et al. (1998). Regulation of interleukin-13 receptor constituents on mature human B lymphocytes. *J Biol Chem*, 273(16), 9864-9871.
- Onoe, Y., Miyaura, C., Kaminakayashiki, T., Nagai, Y., Noguchi, K., Chen, Q. R., et al. (1996). IL-13 and IL-4 inhibit bone resorption by suppressing cyclooxygenase-2-dependent prostaglandin synthesis in osteoblasts. *J Immunol*, 156(2), 758-764.
- Orgogozo, J. M., Gilman, S., Dartigues, J. F., Laurent, B., Puel, M., Kirby, L. C., et al. (2003). Subacute meningoencephalitis in a subset of patients with AD after Abeta42 immunization. *Neurology*, 61(1), 46-54.
- Orr, C. F., Rowe, D. B. and Halliday, G. M. (2002). An inflammatory review of Parkinson's disease. *Progress in Neurobiology*, 68(5), 325-340.
- Patterson, C., Feightner, J. W., Garcia, A., Hsiung, G. Y., MacKnight, C. and Sadovnick, A. D. (2008). Diagnosis and treatment of dementia: 1. Risk assessment and primary prevention of Alzheimer disease. *CMAJ*, 178(5), 548-556.
- Peng, J., Mao, X. O., Stevenson, F. F., Hsu, M. and Andersen, J. K. (2004). The herbicide paraquat induces dopaminergic nigral apoptosis through sustained activation of the JNK pathway. *J Biol Chem*, 279(31), 32626-32632.
- Pollard, H. B., Rojas, E. and Arispe, N. (1993). A new hypothesis for the mechanism of amyloid toxicity, based on the calcium channel activity of amyloid beta protein (A beta P) in phospholipid bilayer membranes. *Ann N Y Acad Sci*, 695, 165-168.
- Pride, M., Seubert, P., Grundman, M., Hagen, M., Eldridge, J. and Black, R. S. (2008). Progress in the active immunotherapeutic approach to Alzheimer's disease: clinical investigations into AN1792-associated meningoencephalitis. *Neurodegener Dis*, 5(3-4), 194-196.
- Punnonen, J. and de Vries, J. E. (1994). IL-13 induces proliferation, Ig isotype switching, and Ig synthesis by immature human fetal B cells. *J Immunol*, 152(3), 1094-1102.
- Rajagopalan, S. and Andersen, J. K. (2001). Alpha synuclein aggregation: is it the toxic gain of function responsible for neurodegeneration in Parkinson's disease? *Mech Ageing Dev*, 122(14), 1499-1510.

- Reiss, A. B. and Wirkowski, E. (2007). Role of HMG-CoA reductase inhibitors in neurological disorders : progress to date. *Drugs*, 67(15), 2111-2120.
- Rideout, H. J., Larsen, K. E., Sulzer, D. and Stefanis, L. (2001). Proteasomal inhibition leads to formation of ubiquitin/alpha-synuclein-immunoreactive inclusions in PC12 cells. *J Neurochem*, 78(4), 899-908.
- Rossi, S., Mancino, R., Bergami, A., Mori, F., Castelli, M., De Chiara, V., et al. (2011). Potential role of IL-13 in neuroprotection and cortical excitability regulation in multiple sclerosis. *Mult Scler*, 17(11), 1301-1312.
- Roy, B., Bhattacharjee, A., Xu, B., Ford, D., Maizel, A. L. and Cathcart, M. K. (2002). IL-13 signal transduction in human monocytes: phosphorylation of receptor components, association with Jaks, and phosphorylation/activation of Stats. *J Leukoc Biol*, 72(3), 580-589.
- Russell, J. A., Ciucci, M. R., Connor, N. P. and Schallert, T. (2010). Targeted exercise therapy for voice and swallow in persons with Parkinson's disease. *Brain Res*, 1341, 3-11.
- Samii, A., Nutt, J. G. and Ransom, B. R. (2004). Parkinson's disease. *Lancet*, 363(9423), 1783-1793.
- Santibanez, M., Bolumar, F. and Garcia, A. M. (2007). Occupational risk factors in Alzheimer's disease: a review assessing the quality of published epidemiological studies. *Occup Environ Med*, 64(11), 723-732.
- Scarmeas, N., Stern, Y., Mayeux, R. and Luchsinger, J. A. (2006). Mediterranean diet, Alzheimer disease, and vascular mediation. *Arch Neurol*, 63(12), 1709-1717.
- Scherzer, C. R., Grass, J. A., Liao, Z., Pepivani, I., Zheng, B., Eklund, A. C., et al. (2008). GATA transcription factors directly regulate the Parkinson's disease-linked gene alpha-synuclein. *Proc Natl Acad Sci U S A*, 105(31), 10907-10912.
- Schindler, C., Fu, X. Y., Improta, T., Aebbersold, R. and Darnell, J. E., Jr. (1992). Proteins of transcription factor ISGF-3: one gene encodes the 91- and 84-kDa ISGF-3 proteins that are activated by interferon alpha. *Proc Natl Acad Sci U S A*, 89(16), 7836-7839.

- Schoenly, K. A. and Weiner, D. B. (2008). Human immunodeficiency virus type 1 vaccine development: recent advances in the cytotoxic T-lymphocyte platform "spotty business". *J Virol*, 82(7), 3166-3180.
- Shavali, S., Combs, C. K. and Ebadi, M. (2006). Reactive macrophages increase oxidative stress and alpha-synuclein nitration during death of dopaminergic neuronal cells in co-culture: relevance to Parkinson's disease. *Neurochem Res*, 31(1), 85-94.
- Silvennoinen, O., Ihle, J. N., Schlessinger, J. and Levy, D. E. (1993). Interferon-induced nuclear signalling by Jak protein tyrosine kinases. *Nature*, 366(6455), 583-585.
- Spillantini, M. G., Divane, A. and Goedert, M. (1995). Assignment of human alpha-synuclein (SNCA) and beta-synuclein (SNCB) genes to chromosomes 4q21 and 5q35. *Genomics*, 27(2), 379-381.
- Szekely, C. A., Breitner, J. C., Fitzpatrick, A. L., Rea, T. D., Psaty, B. M., Kuller, L. H., et al. (2008). NSAID use and dementia risk in the Cardiovascular Health Study: role of APOE and NSAID type. *Neurology*, 70(1), 17-24.
- Szekely, C. A., Breitner, J. C. and Zandi, P. P. (2007). Prevention of Alzheimer's disease. *Int Rev Psychiatry*, 19(6), 693-706.
- Tanner, J. W., Chen, W., Young, R. L., Longmore, G. D. and Shaw, A. S. (1995). The conserved box 1 motif of cytokine receptors is required for association with JAK kinases. *J Biol Chem*, 270(12), 6523-6530.
- Thiruchelvam, M., McCormack, A., Richfield, E. K., Baggs, R. B., Tank, A. W., Di Monte, D. A., et al. (2003). Age-related irreversible progressive nigrostriatal dopaminergic neurotoxicity in the paraquat and maneb model of the Parkinson's disease phenotype. *European Journal of Neuroscience*, 18(3), 589-600.
- Thompson, C. A., Spilsbury, K., Hall, J., Birks, Y., Barnes, C. and Adamson, J. (2007). Systematic review of information and support interventions for caregivers of people with dementia. *BMC Geriatr*, 7, 18.
- Tiraboschi, P., Hansen, L. A., Thal, L. J. and Corey-Bloom, J. (2004). The importance of neuritic plaques and tangles to the development and evolution of AD. *Neurology*, 62(11), 1984-1989.

- Uversky, V. N., Li, J. and Fink, A. L. (2001). Pesticides directly accelerate the rate of alpha-synuclein fibril formation: a possible factor in Parkinson's disease. *FEBS Lett*, 500(3), 105-108.
- Van Broeck, B., Van Broeckhoven, C. and Kumar-Singh, S. (2007). Current insights into molecular mechanisms of Alzheimer disease and their implications for therapeutic approaches. *Neurodegener Dis*, 4(5), 349-365.
- Veldman, B. A., Wijn, A. M., Knoers, N., Praamstra, P. and Horstink, M. W. (1998). Genetic and environmental risk factors in Parkinson's disease. *Clin Neurol Neurosurg*, 100(1), 15-26.
- Vila, M., Vukosavic, S., Jackson-Lewis, V., Neystat, M., Jakowec, M. and Przedborski, S. (2000). Alpha-synuclein up-regulation in substantia nigra dopaminergic neurons following administration of the parkinsonian toxin MPTP. *J Neurochem*, 74(2), 721-729.
- von Rotz, R. C., Kohli, B. M., Bosset, J., Meier, M., Suzuki, T., Nitsch, R. M., et al. (2004). The APP intracellular domain forms nuclear multiprotein complexes and regulates the transcription of its own precursor. *J Cell Sci*, 117(19), 4435-4448.
- Waldemar, G., Dubois, B., Emre, M., Georges, J., McKeith, I. G., Rossor, M., et al. (2007). Recommendations for the diagnosis and management of Alzheimer's disease and other disorders associated with dementia: EFNS guideline. *Eur J Neurol*, 14(1), e1-26.
- Wang, Y., Zhang, Y., Wei, Z., Li, H., Zhou, H. and Zhang, Z. (2009). JNK inhibitor protects dopaminergic neurons by reducing COX-2 expression in the MPTP mouse model of subacute Parkinson's disease. *J Neurol Sci*, 285(1-2), 172-177.
- Waring, S. C. and Rosenberg, R. N. (2008). Genome-wide association studies in Alzheimer disease. *Arch Neurol*, 65(3), 329-334.
- Whitton, P. S. (2007). Inflammation as a causative factor in the aetiology of Parkinson's disease. *Br J Pharmacol*, 150(8), 963-976.

- Wichmann, T. and DeLong, M. R. (2007). Anatomy and physiology of the basal ganglia: Relevance to Parkinson's disease and related disorders. *Handbook of Clinical Neurology*, 83, 1-18.
- Withers, G. S., George, J. M., Banker, G. A. and Clayton, D. F. (1997). Delayed localization of synelfin (synuclein, NACP) to presynaptic terminals in cultured rat hippocampal neurons. *Brain Res Dev Brain Res*, 99(1), 87-94.
- World Health Organization. (2006). *Neurological disorders: Public health challenges*. Switzerland: World Health Organization.
- Wu, X. F., Block, M. L., Zhang, W., Qin, L., Wilson, B., Zhang, W. Q., et al. (2005). The role of microglia in paraquat-induced dopaminergic neurotoxicity. *Antioxid Redox Signal*, 7(5-6), 654-661.
- Xia, Y., Saitoh, T., Ueda, K., Tanaka, S., Chen, X., Hashimoto, M., et al. (2001). Characterization of the human alpha-synuclein gene: Genomic structure, transcription start site, promoter region and polymorphisms. *J Alzheimers Dis*, 3(5), 485-494.
- Xing, X., Sha, S., Li, Y., Zong, L., Jiang, T. and Cao, Y. (2012). Immunization with a new DNA vaccine for Alzheimer's disease elicited Th2 immune response in BALB/c mice by in vivo electroporation. *J Neurol Sci*, 313(1-2), 17-21.
- Yang, W., Tiffany-Castiglioni, E., Koh, H. C. and Son, I. H. (2009). Paraquat activates the IRE1/ASK1/JNK cascade associated with apoptosis in human neuroblastoma SH-SY5Y cells. *Toxicol Lett*, 191(2-3), 203-210.
- Yang, W., Tiffany-Castiglioni, E., Lee, M. Y. and Son, I. H. (2010). Paraquat induces cyclooxygenase-2 (COX-2) implicated toxicity in human neuroblastoma SH-SY5Y cells. *Toxicol Lett*, 199(3), 239-246.
- Yu, M. S., Suen, K. C., Kwok, N. S., So, K. F., Hugon, J. and Chang, R. C. (2006). Beta-amyloid peptides induces neuronal apoptosis via a mechanism independent of unfolded protein responses. *Apoptosis*, 11(5), 687-700.

Zhong, Z., Wen, Z. and Darnell, J. E., Jr. (1994). Stat3: a STAT family member activated by tyrosine phosphorylation in response to epidermal growth factor and interleukin-6. *Science*, 264(5155), 95-98.

Zhu, C., Zhang, A., Huang, S., Ding, G., Pan, X. and Chen, R. (2010). Interleukin-13 inhibits cytokines synthesis by blocking nuclear factor- κ B and c-Jun N-terminal kinase in human mesangial cells. *J Biomed Res*, 24(4), 308-316.

