



REFERENCES

มหาวิทยาลัยนเรศวร

REFERENCES

- Agrawal, N., Dasaradhi, P. V. N., Mohammed, A., Malhotra, P., Bhatnagar, R. K. and Mukherjee, S. K. (2003). RNA Interference: Biology, Mechanism, and Applications. **Microbiology and Molecular Biology Reviews**, 67(4), 657-685.
- Aida, R., Kishimoto, S., Tanaka, Y. and Shibata, M. (2000). Modification of flower color in torenia (*Torenia fournieri* Lind.) by genetic transformation. **Plant Science**, 153(1), 33-42.
- Barampuram, S. and Zhang, Z. J. (2011). Recent advances in plant transformation. **Methods in Molecular Biology**, 701, 1-35.
- Brugliera, F., Tull, D., Holton, T. A., Karan, M., Treloar, N., Simpson, K., et al. (2000). Introduction of a cytochrome b5 enhances the activity of flavonoid 3',5'-hydroxylase (a cytochrome P450) in transgenic carnation. **International Plant Molecular Biology Reporter**, 18(2), S6-S8.
- Finer, J. J. (2010). Plant Nuclear Transformation. **Biotechnology in Agriculture and Forestry**, 64, 3-21.
- Forkmann, G. and Martens, S. (2001). Metabolic engineering and applications of flavonoids. **Current Opinion in Biotechnology**, 12(2), 155-160.
- Gil-Humanes, J., Pistón, F., Hernando, A., Alvarez, J. B., Shewry, P. R. and Barro, F. (2008). Silencing of γ -gliadins by RNA interference (RNAi) in bread wheat. **Cereal Science**, 48(3), 565-568.
- Goodin, M. M., Zaitlin, D., Naidu, R. A. and Lommel, S. A. (2008). *Nicotiana benthamiana*: its history and future as a model for plant-pathogen interactions. **Molecular Plant-Microbe Interactions**, 21(8), 1015-1026.
- Hamada, W. and Spanu, P. D. (1998). Co-suppression of the hydrophobin gene HCf-1 is correlated with antisense RNA biosynthesis in *Cladosporium fulvum*. **Molecular and General Genetics**, 259(6), 630-638.
- Hoffmann, T., Kalinowski, G. and Schwab, W. (2006). RNAi-induced silencing of gene expression in strawberry fruit (*Fragaria x ananassa*) by agroinfiltration: a rapid assay for gene function analysis. **Plant Journal**, 48(5), 818-826.

- Holton, T. A. and Cornish, E. C. (1995). Genetics and Biochemistry of Anthocyanin Biosynthesis. *Plant Cell*, 7(7), 1071-1083.
- Hung, K. T., Cheng, D. G., Hsu, Y. T. and Kao, C. H. (2008). Abscisic acid-induced hydrogen peroxide is required for anthocyanin accumulation in leaves of rice seedlings. *Plant Physiology*, 165(12), 1280-1287.
- Jiang, F., Wang, J.-Y., Jia, H.-F., Jia, W.-S., Wang, H.-Q. and Xiao, M. (2013). RNAi-Mediated Silencing of the Flavanone 3-Hydroxylase Gene and Its Effect on Flavonoid Biosynthesis in Strawberry Fruit. *Plant Growth Regulation*, 32(1), 182-190.
- Jones, A. L., Thomas, C. L. and Maule, A. J. (1998). *De novo* methylation and co-suppression induced by a cytoplasmically replicating plant RNA virus. *EMBO Journal*, 17(21), 6385-6393.
- Jones, H. D., Doherty, A. and Sparks, C. A. (2009). Transient transformation of plants. *Methods in Molecular Biology*, 513, 131-152.
- Jones, H. D. and Sparks, C. A. (2009). Stable transformation of plants. *Methods in Molecular Biology*, 513, 111-130.
- Kamiishi, Y., Otani, M., Takagi, H., Han, D.-S., Mori, S., Tatsuzawa, F., et al. (2012). Flower color alteration in the liliaceous ornamental *Tricyrtis* sp. by RNA interference-mediated suppression of the chalcone synthase gene. *Molecular Breeding*, 30(2), 671-680.
- Katsumoto, Y., Fukuchi-Mizutani, M., Fukui, Y., Brugliera, F., Holton, T. A., Karan, M., et al. (2007). Engineering of the rose flavonoid biosynthetic pathway successfully generated blue-hued flowers accumulating delphinidin. *Plant Cell Physiology*, 48(11), 1589-1600.
- Koseki, M., Goto, K., Masuta, C. and Kanazawa, A. (2005). The star-type color pattern in *Petunia hybrida* 'red Star' flowers is induced by sequence-specific degradation of chalcone synthase RNA. *Plant Cell Physiology*, 46(11), 1879-1883.
- Kubo, A., Aono, M., Nakajima, N., Saji, H., Tanaka, K. and Kondo, N. (1999). Differential Responses in Activity of Antioxidant Enzymes to Different Environmental Stresses in *Arabidopsis thaliana*. *Plant Research*, 112(3), 279-290.

- Kunu, W., Thanonkeo, S. and Thanonkeo, P. (2012). Cloning and expression analysis of dihydroxyflavonol 4-reductase (*DFR*) in *Ascocenda* spp. **Biotechnology**, 11(64), 12702-12709.
- Liew, C.-F., Loh, C.-S., Goh, C.-J. and Lim, S.-H. (1998). The isolation, molecular characterization and expression of dihydroflavonol 4-reductase cDNA in the orchid, *Bromheadia finlaysoniana*. **Plant Science**, 135(2), 161-169.
- Ma, H. and Pooler, M. (2009). Anthocyanin Regulatory/Structural Gene Expression in *Phalaenopsis*. **the American Society for Horticultural Science**, 134(1), 88-96.
- Marina biotech. (2013). **RNA interference**. Retrieved May 18, 2013, from http://www.marinabio.com/rna_interference.
- Meyer, P., Heidmann, I., Forkmann, G. and Saedler, H. (1987). A new petunia flower color generated by transformation of a mutant with a maize gene. **Nature**, 330(6149), 677-678.
- Monmai, C. and Ratanasut, K. (2012). An improved method for RNA Miniprep from flowers of *Dendrobium* Sonia Earsakul for Northern analysis. **Proceeding: 1st ASEAN plus three graduate research congress**, 1, ST-412-416.
- Mudalige-Jayawickrama, R. G., Champagne, M. M., Hieber, A. D. and Kuehnle, A. R. (2005). Cloning and Characterization of Two Anthocyanin Biosynthetic Genes from *Dendrobium* Orchid. **the American Society for Horticultural Science**, 130(4), 611-618.
- Nakatsuka, T., Abe, Y., Kakizaki, Y., Yamamura, S. and Nishihara, M. (2007). Production of red-flowered plants by genetic engineering of multiple flavonoid biosynthetic genes. **Plant Cell Reports**, 26(11), 1951-1959.
- Nakatsuka, T., Mishiba, K.-i., Abe, Y., Kubota, A., Kakizaki, Y., Yamamura, S., et al. (2008). Flower color modification of gentian plants by RNAi-mediated gene silencing. **Plant Biotechnology**, 25(1), 61-68.

- Nakatsuka, T., Mishiba, K.-i., Kubota, A., Abe, Y., Yamamura, S., Nakamura, N., et al. (2010). Genetic engineering of novel flower color by suppression of anthocyanin modification genes in gentian. *Plant Physiology*, 167(3), 231-237.
- Nakatsuka, T., Nishihara, M., Mishiba, K. and Yamamura, S. (2005). Temporal expression of flavonoid biosynthesis-related genes regulates flower pigmentation in gentian plants. *Plant Science*, 168(5), 1309-1318.
- Newell, C. (2000). Plant transformation technology. *Molecular Biotechnology*, 16(1), 53-65.
- Nielsen, K., Deroles, S., Markham, K., Bradley, M., Podivinsky, E. and Manson, D. (2002). Antisense flavonol synthase alters copigmentation and flower color in lisianthus. *Molecular Breeding*, 9(4), 217-229.
- Nishihara, M., Nakatsuka, T., Mishiba, K., Kikuchi, A. and Yamamura, S. (2003). Flower Color Modification by Suppression of Chalcone Synthase Gene in Gentian. *Plant Cell Physiology*, 44, S159.
- Office of Agricultural Economics. (2013). Agricultural import export. Retrieved May 18, 2013, from http://www.oae.go.th/oaе_report/export_import/export_result.php
- Orzaez, D., Mirabel, S., Wieland, W. H. and Granell, A. (2006). Agroinjection of tomato fruits. A tool for rapid functional analysis of transgenes directly in fruit. *Plant Physiology*, 140(1), 3-11.
- Park, J.-S., Choung, M.-G., Kim, J.-B., Hahn, B.-S., Kim, J.-B., Bae, S.-C., et al. (2007). Genes up-regulated during red coloration in UV-B irradiated lettuce leaves. *Plant Cell Reports*, 26(4), 507-516.
- Pitakdantham, W., Sutabutra, T., Chiemsombat, P. and Pitaksutheepong, C. (2011). Isolation and Characterization of Dihydroflavonol 4-reductase Gene in *Dendrobium* Flowers. *Plant Sciences*, 6(2), 88-94.
- Polashock, J. J., Griesbach, R. J., Sullivan, R. F. and Vorsa, N. (2002). Cloning of a cDNA encoding the cranberry dihydroflavonol-4-reductase (DFR) and expression in transgenic tobacco. *Plant Science*, 163(2), 241-251.

- Reilly, C. O., Shepherd, N. S., Pereira, A., Schwarz-Sommer, Z. and Bertram, I. (1985). Molecular cloning of the a7 locus in Zea mays using the transposable elements En and Mu7. **EMBO Journal**, 4, 877-882.
- Royal horticultural societies. (2011). **Orchids**. Retrieved May 21, 2013, from <http://www.rhs.org.uk/Plants/Plant-science/Plant-registration/Orchids>.
- Saito, R., Fukuta, N., Ohmiya, A., Itoh, Y., Ozeki, Y., Kuchitsu, K., et al. (2006). Regulation of anthocyanin biosynthesis involved in the formation of marginal picotee petals in *Petunia*. **Plant Science**, 170(4), 828-834.
- Santos-Rosa, M., Poutaraud, A., Merdinoglu, D. and Mestre, P. (2008). Development of a transient expression system in grapevine via agro-infiltration. **Plant Cell Reports**, 27(6), 1053-1063.
- Shang, Y., Schwinn, K., Bennett, M., Hunter, D., Waugh, T., Pathirana, N., et al. (2007). Methods for transient assay of gene function in floral tissues. **Plant Methods**, 3(1), 1-12.
- Sparkes, I. A., Runions, J., Kearns, A. and Hawes, C. (2006). Rapid, transient expression of fluorescent fusion proteins in tobacco plants and generation of stably transformed plants. **NAT Protocol Journal**, 1(4), 2019-2025.
- Suzuki, K.-i., Xue, H.-m., Tanaka, Y., Fukui, Y., Fukuchi-Mizutani, M., Murakami, Y., et al. (2000). Flower color modifications of *Torenia hybrida* by cosuppression of anthocyanin biosynthesis genes. **Molecular Breeding**, 6(3), 239-246.
- Tanaka, Y., Brugliera, F., Kalc, G., Senior, M., Dyson, B., Nakamura, N., et al. (2010). Flower color modification by engineering of the flavonoid biosynthetic pathway: practical perspectives. **Bioscience, Biotechnology, and Biochemistry**, 74(9), 1760-1769.
- Tanaka, Y., Tsuda, S. and Kusumi, T. (1998). Metabolic Engineering to Modify Flower Color. **Plant Cell Physiology**, 39(11), 1119-1126.
- To, K.-Y. and Wang, C.-K. (2006). **Molecular Breeding of Flower Color**. Floriculture, Ornamental and Plant Biotechnology. UK: Global Science Books.

- Tsai, W.-C., Hsiao, Y.-Y., Lee, S.-H., Tung, C.-W., Wang, D.-P., Wang, H.-C., et al. (2006). Expression analysis of the ESTs derived from the flower buds of *Phalaenopsis equestris*. *Plant Science*, 170(3), 426-432.
- Tsuda, S., Fukui, Y., Nakamura, N., Katsumoto, Y., Yonekura-Sakakibara, K., Fukuchi-Mizutani, M., et al. (2004). Flower color modification of *Petunia hybrida* commercial varieties by metabolic engineering. *Plant Biotechnology*, 21(5), 377–386.
- Ueyama, Y., Katsumoto, Y., Fukui, Y., Fukuchi-Mizutani, M., Ohkawa, H., Kusumi, T., et al. (2006). Molecular characterization of the flavonoid biosynthetic pathway and flower color modification of *Nierembergia* sp. *Plant Biotechnology*, 23, 19–24.
- Ueyama, Y., Suzuki, K.-i., Fukuchi-Mizutani, M., Fukui, Y., Miyazaki, K., Ohkawa, H., et al. (2002). Molecular and biochemical characterization of torenia flavonoid 3'-hydroxylase and flavone synthase II and modification of flower color by modulating the expression of these genes. *Plant Science*, 163(2), 253-263.
- Van Haute, E., Joos, H., Maes, M., Warren, G., Van Montagu, M. and Schell, J. (1983). Intergeneric transfer and exchange recombination of restriction fragments cloned in pBR322: a novel strategy for the reversed genetics of the Ti plasmids of *Agrobacterium tumefaciens*. *EMBO Journal*, 2(3), 411-417.
- Wroblewski, T., Tomczak, A. and Michelmore, R. (2005). Optimization of Agrobacterium-mediated transient assays of gene expression in lettuce, tomato and Arabidopsis. *Plant Biotechnol*, 3(2), 259-273.
- Yang, Y., Li, R. and Qi, M. (2000). *In vivo* analysis of plant promoters and transcription factors by agroinfiltration of tobacco leaves. *Plant Journal*, 22(6), 543-551.
- Yasmin, A. and Debener, T. (2010). Transient gene expression in rose petals via *Agrobacterium* infiltration. *Plant Cell, Tissue and Organ Culture* 102(2), 245-250.
- Zhang, P., Wen, P.-F., Wan, S.-B., Wang, W., Pan, Q.-H., Zhan, J.-C., et al. (2008). Molecular cloning of dihydroflavonol 4-reductase gene from grape berry and preparation of an anti-DFR polyclonal antibody. *Plant Vitis*, 47(3), 141–145.

Zottini, M., Barizza, E., Costa, A., Formentin, E., Ruberti, C., Carimi, F., et al. (2008). Agroinfiltration of grapevine leaves for fast transient assays of gene expression and for long-term production of stable transformed cells. *Plant Cell Reports*, 27(5), 845-853.

