

REFERENCES



REFERENCES

- Alvarez-Roman R. et al. (2004). Visualization of skin panatration using confocal laser scanning microscopy. Eur. J. Pharm. Biopharm, 58, 301-316.
- Athanasios S. et al. (2006). Phathogenic role for skin macrophages in a mouse model of keratinocyte induced psoriasis like skin inflammation. J.Am.Aca.Dermatol, 116(8), 2094-2104.
- Avinash K.S. & Ambikanandan M. (2002). Mathmetical modeling of preparation of acyclovir liposomes: reverse phase evaporation method. J. Pharm. Pharmaceut. Sci, 5(3), 285-291.
- Bangham A.D. et al. (1965). Diffusion of univalent ions across the lamellar of swollen phospholipids. J. Mol. Biol, 13, 238-252.
- Bergfield W.F. et al. (1997). Cosmetic use of alpha hydroxyl acids. Cleveland Clin.J.Med, 64, 327-329.
- Betz G. et al. (2001). Heparin penetration into and permeation through human skin from aqueous and liposomal formulation in vitro. Int. J. Pharm, 228, 147-159.
- Betz G. et al. (2001). Interaction of liposome formulations with human skin in vitro. Int. J. Pharm, 229, 117-129.
- Betz G. et al. (2005). In vivo comparison of various liposome formulations for cosmetic application. Int. J. Pharm, 296, 44-54.
- Bolognia, J.L. & Orlow, S.J. (1999). Melanocyte biology. Retrieved on May 14, 2002.
- Coderch L. et al. (1996). Percutaneous penetration of liposomes using the tape stripping technique. Int. J. Pharm, 139, 197-203.
- Copovi A. et al. (2006). Enhancing effect of AHAs on in vitro permeation across the human skin of compounds with different lipophilicity. Int. J. Pharm, 314, 31-36.
- Contreras F. et al. (2005). In vitro percutaneous absorption of all-trans retinoic acid applied in free from or encapsulated in stratum corneum lipid liposomes. Int. J. Pharm, 297, 134-145.

- Dayong Z. Et al. (2001). Leakage and aggregation of phospholipid vesicles induced by the BH3-only Bcl-2 family member, BID, 286, 48-55.
- De Maria P. et al. (2005). Cardanol as a replacement for cholesterol into lipid bilayer of POPC liposomes. Colloids and Surface B, 40, 11-18.
- Dassanayake M. D. & Fosberg, F. R. (Eds.). (1991). A Revised Handbook to the Flora of Ceylon. Washington, D. C.: ASM Press.
- Ditre C.M. (1996). Effect of alpha hydroxyl acids on photoaged skin: a pilot clinical, histologic and ultrastructural study. J. Am. Aca. Dermatol, 34, 187-195.
- Feng S., Ruan G., Li Q. (2004). Fabrication and characterizations of a novel drug delivery device liposomes in microsphere. Biomaterials, 25, 5181-5189.
- Elsner, P. & Maibach H.I. (Eds). (2002). Depigmentation agents. New York: Marcel Dekker.
- Freedberg, I.M. & Fitzpatrick, T.B. (Eds). (1993). Biology of melanocytes New York: MacGraw-Hill.
- Gomez-Hens A. & Fernandez-Romero J. M. (2005). The role of liposomes in analytical processes. Trend. Anal. Chem, 24(1), 9-19.
- Guo J. et al. (2003). Chitosan coated liposomes: characterization and interaction with leoprolide. Int.J.Pharm, 260, 167-173.
- Hwang M. et al. (2001). Stabilization of phosphatidylserine/ phosphatidylethanolamine liposomes with hydrophilic polymers having multiple sticky feet. J. Sur. Coll, 17(25), 7713-7716.
- Jiang M. & Qureshi S.A. (1998). Assessment of in vitro percutaneous absorption of glycolic acid through human skin section using a flow-through diffusion cell system. J. Dermatol. Sci, 18, 181-188.
- Jin Ho C. et al. (2003). Aging and photoaging. J. Am. Acad. Dermatol, 49(4), 690-697.
- Lopez O. et al. (2002). Reconstitution of liposomes inside the intercellular lipid domain of the stratum corneum. Colloid and Surfaces, 18, 7002-7008.
- Males R.G. & Herring F.G. (1999). A H-NMR study of the permeation of glycolic acid through phospholipid membrane. Biochimica et Biophysica Acta, 1416, 333-338.

- Maria B. et al. (2001). Stratum corneum lipids liposomes for the topical delivery of 5-aminolevulinic acid in photodynamic therapy of skin cancer: preparation and in vitro permeation study. BMC Dermatology, 1471, 1-5.
- Maria P. et al. (2005). Cardanol as a replacement for cholesterol into the lipid bilayer of POPC liposome. Colloid and Surfaces, 40, 11-18.
- Mats A. et al. (2000). Cryo transmission electron microscopy of liposomes and related structure. Colloid. Sur, 174, 3-21.
- McPhail D. et al. (2000). Liposomes encapsulating polymeric chitosan based vesicles-a vehicle in vesicle system for drug delivery. Int. J. Pharm, 200, 73-86.
- Nounou M.M. et al. (2006). In vitro release of hydrophilic and hydrophobic drugs from liposomal and gels. Acta. Pharm, 56, 311-324.
- Olga L. et al. (2002). Reconstitution of liposomes inside the intercellular lipid domain of the stratum corneum. Langmuir, 18, 7002-7008.
- Omar M. et al. (2005). Production of soybean phosphatidylcholine-chitosan nanovesicles by reverse phase evaporation. Chem. Phys. Lip, 138, 29-37.
- Perugini P. et al. (2000). Study on glycolic acid delivery by liposomes and microspheres. Int. J. Pharm, 196, 51-61.
- Ponec, M. (2002). Skin constructs for replacement skin tissue for *in vitro* testing. Advanced Drug Delivery, 54 (1), 19-30
- Riccardo A. & Muzzarelli A. (1998). Colorimetric determination of chitosan. Anal. Biochem, 260, 255-257.
- Ruel-Gariepy E. et al. (2002). Thermosensitive chitosan-bases hydrogel containing liposomes for the delivery of hydrophilic molecules. J. Control. Release, 103, 123-136.
- Seiberg, M. (2001). Keratinocyte-melanocyte interactions during melanosome transfer. Pigment Cell Research, 14 (4), 236-242.
- Shinsuke O. et al. (2006). Prepration of double liposomes and their efficiency as an oral vaccine carrier. Biol. Pharm. Bull, 29(6), 1223-1228.
- Shuji K. & Masatshi K. (2006). Enhance delivery of retinoic acid to skin by cationic liposomes. Chem. Pharm. Bull, 54(2), 242-244.

- Sinico C. et al. (2005). Liposomes as carriers for dermal delivery of tretinoin: in vitro evaluation of drug permeation and vesicle-skin interaction. J. Control. Release, 103, 123-136.
- Sulkowski W.W. et al. (2005). The influence of temperature, cholesterol content and pH on liposome stability. J. Mol. Struct., 25, 1-11
- Sumio C. et al. (2006). Pharmacokinetic analysis of the uptake of liposomes by macrophages and foam cells in vitro and their distribution to atherosclerotic lesions in mice. J. Control. Release, 21(1), 37-44.
- Takeuchi H. et al. (2003). Muscoadhesive properties of carbopol or chitosan coated. J. Control. Release, 86, 235-242.
- Takeuchi H. et al. (2005). Effectiveness of submicron sized chitosan coated liposomes in oral administration of peptide drugs. Int. J. Pharm, 303, 160-170.
- Tsen-Fang T. et al. (2000). Effect of glycolic acid on light-induced skin pigmentation in Asian and Caucasian subjects. J. Am. Acad. Dermatol, 43(2), 238-243.
- Ulf M. et al. (2004). Human leukocyte elastase induces keratinocyte proliferation by epidermal growth factor receptor activation. J. Invest. Dermatol, 123(2), 338-345.
- Usuki A. et al. (2003). The inhibitory effect of glycolic acid and lactic acid on melanin synthesis in melanoma cells. Exp. Dermatol, 12(2), 43-50.
- Van Scott E.J. et al. (1996). Alpha hydroxyacids in the treatment of sign of photoaging, Clin. Dermatol, 14, 217-226.
- Vermuri S. & Rhodes C.T. (1995). Developmaent and validation of a drug release rate method for a water soluble drug in a liposome preparation. Drug. Dev. Ind. Pharm, 70, 95-111.
- Viyoch J. et al. (2003). Formulation and development of a patch containing tamarind fruit extract by using the blended rchitosan-starch as a rate controlling matrix. Int. J. Cos. Sci, 25, 113-125.
- Wan Y. et al. (2002). Liposomes from novel photolabile:light-induced unloaded of small molecules as monitored by PFG NMR. J. Am. Chem. Soc, 124, 5610-5611.

- Weiner N., Martin F. & Riaz M. (1989). Liposomes as a drug delivery system. Drug. Develop. Indust. Pharm, 15(10),1523-1551.
- Walters, K.A (Ed). 2002. The structure and function of skin. New York: Marcel Dekker.
- Woodle M.C. & Lasic D.D. (1992). Sterically stabilized liposomes. Biochim. Biophys. Acta, 113, 171-199.
- Yamamoto Y. et al. (2006). Effects of alpha-hydroxy acids on the human skin of Japanese subjects. J. Dermatol, 33(1), 16-22.
- Yasutami S. et al. (2004). Suppressed permeation of linoleic acid in a liposomal formulation through reconstructed skin tissue. Biol. Pharm. Bull, 27(6), 879-882.
- Yongqin W. Et al. (2002). Liposomes from novel photolabile phospholipids. J. Am. Chem. Soc, 124, 5610-5611.
- Zeng-hong W. et al. (2004). Hypoglycemic efficacy of chitosan-coated insulin liposomes after oral administration in mice. Pharmacol. Sin, 25(7), 996-972.
- Zygmunt H.Z. (2004). Vesicles with a double bilayer. Cell. Mol. Biol. Lett, 9(4), 589-602.