



REFERENCES

มหาวิทยาลัยพระศรี

REFERENCES

- Adebowale, K.O. et al. (2005). Functional properties of native, physically and chemically modified breadfruit (*Artocarpus artilis*) starch. Industrial Crops and Products, 21 (3), 343-351.
- Amarowicz, R. et al. (2004). Free-radical scavenging capacity and antioxidant activity of selected plant species from the Canadian prairies. Food Chemistry, 85 (4), 551-562.
- Amusa, N.A., Kehinde, I. & Ashaye, O. (2002). Bio-deterioration of breadfruit (*Artocarpus Communis*) in storage and its effects on the nutrient composition. African Journal of Biotechnology, 1 (2), 57-60.
- Ancans, J. & Thody, A.J. (2000). Activation of melanogenesis by vacuolar type H^+ -ATPase inhibitors in amelanotic, tyrosinase positive human and mouse melanoma cells. FEBS Letters, 478 (1-2), 57-60.
- Bologna, J.L. & Orlow, S.J. (November 6, 1999). Melanocyte biology. Retrieved on May 14, 2002 from: <http://www3.us.elsevierhealth.com/HS/promo/Bologna/Ch65.pdf>.
- Briganti, S., Camera, E. & Picardo, M. (2003). Chemical and instrumental approaches to treat hyperpigmentation. Pigment Cell Research, 16 (2), 101-110.
- Cevc, G. (2004). Lipid vesicles and other colloids as drug carriers on the skin. Advanced Drug Delivery, 56 (5), 675-711.
- Chang, M.L. & Chang, C.M. (2003). Simultaneous HPLC determination of hydrophilic whitening agents in cosmetic products. Journal of Pharmaceutical and Biomedical Analysis, 33 (4), 617-626.
- Choi, S.Y. et al. (2002). (4-methoxy-benzylidene)-(3-methoxy-phenyl)-amine, a nitrogen analog of stilbene as a potent inhibitor of melanin production. Chemical & Pharmaceutical Bulletin, 50 (4), 450-452.
- Curto, E.V. et al. (1999). Inhibitor of mammalian melanocyte tyrosinase: *in vitro* comparisons of alkyl esters of gentisic acid with others putative inhibitors. Biochemical Pharmacology, 57 (6), 663-672.

- de Leeuw, S.M. et al. (2001). Melanin content of cultured human melanocytes and UV-induced cytotoxicity. Journal of Photochemistry and Photobiology, 61 (3), 106-113.
- Elsner, P. & Maibach H.I. (Eds). (2002). Depigmentation agents. New York: Marcel Dekker.
- Eves, P.C., MacNeil, S. & Haycock, J.W. (2006). α -Melanocyte stimulating hormone, inflammation and human melanoma. Peptides, 27 (2), 444-452.
- Freedberg, I.M. & Fitzpatrick, T.B. (Eds). (1993). Biology of melanocytes New York. MacGraw-Hill.
- Hearing, V.J. & Tsukamoto K. (1991). Enzymatic control of pigmentation in mammals. The FASEB Journal, 5 (14), 290-299.
- Hearing, V.J. (2005). Biogenesis of pigment granules: a sensitive way to regulate melanocyte function. Journal of Dermatological Science, 37 (1), 3-14.
- Huang, K.F. et al. (2005). Studies on the inhibitory effect of *Graptotalum paraguayense* E. Walter extract on mushroom tyrosinase. Food Chemistry, 89 (4), 583-587.
- Juliano, C. et al. (2005). Antioxidant activity of gamma-oryzanol: mechanism of action and its effect on oxidative stability of pharmaceutical oils. International Journal of Pharmaceutics, 299 (1-2), 146-154.
- Jung, G.D. et al. (2001). Stimulation of melanogenesis by glycyrrhizin in B16 melanoma cells. Experimental and Molecular Medicine, 33 (3), 131-135.
- Katsambas, A.D. & Stratigos, A.J. (2001). Depigmenting and bleaching agents: coping with hyperpigmentation. Clinic in Dermatology, 191 (4), 483-488.
- Khatib, S. et al. (2005). Chalcones as potent tyrosinase inhibitors: the importance of a 2, 4-substitute resorcinol moiety. Bioorganic & Medicinal Chemistry, 13 (2), 433-441.
- Kim, D.H. et al. (2003). Development of 5-[(3-aminopropyl) phosphinoxy]-2-(hydroxymethyl)-4H-pyran-4-one as a novel whitening agent. Chemical & Pharmaceutical Bulletin, 51 (2), 113-116.

- Kim, Y.M. et al. (2002). Oxyresveratrol and hydroxystilbene compounds; inhibitory effect on tyrosinase and mechanism of action. Journal of Biological Chemistry, 277 (18), 16340-16344.
- Kubo, I., Nihei, K. & Tsujimoto, K. (2004). Methyl p- coumarate, a melanin formation inhibitor in B16 mouse melanoma cells. Bioorganic & Medicinal Chemistry, 12 (20), 5349-5354.
- Lee, H.S. (2002). Tyrosinase inhibitors of *Pulsatilla cernua* root-derived materials. Journal of Agriculture Food Chemistry, 50 (6), 1400-1403.
- Lee, J., Koo, N. & Min, D.B. (2003). Reactive oxygen species, aging, and antioxidative nutraceuticals. Institute of Food Technologists, 3 (1), 21-32.
- Lee, S.H. et. al. (2002). Mulberroside F isolated from the leaves of *Morus alba* inhibits melanin biosynthesis. Biological and Pharmaceutical Bulletin, 25 (8), 1045-1408.
- Lei, T.C. et al. (2002). A melanocyte-keratinocyte coculture model to assess regulators of pigmentation in vitro. Analytical Chemistry, 305 (2), 260-268.
- Likhitwitayawuid, K. & Sritularak, B. (2001). A new dimeric stilbene with tyrosinase inhibitory activity from *Artocarpus gomezianus*. Journal of Natural Products, 64 (11), 1457-1459.
- Maeda, K. et al. (1996). Comparison of the melanogenesis in human black and light brown melanocytes. Journal of Dermatological Science, 14 (3), 199-206.
- Mallick, S., Mondal, S.K. & Bhadra, R. (2002). Human placental lipid induces mitogenesis and melanogenesis in B16F10 melanoma cells. Journal of Bioscience, 27 (3), 243-249.
- Mambro, V.M. & Fonseca, M.J.V. (2005). Assays of physical stability and antioxidant activity of a topical formulation added with different plant extracts. Journal of Pharmaceutical and Biomedical Analysis, 37 (2), 287-295.
- Masamoto, Y., Iida, S. & Kubo, M. (1980). Inhibitory effect of Chinese crude drugs on tyrosinase. Planta Medica, 40 (4), 361-365.

- Molyneux, P. (2004). The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity. Songklanakarin Journal of Science and Technology, 26 (2), 211-219.
- Mun, Y.J. et al. (2004). Inhibitory effect of miconazole on melanogenesis. Biological and Pharmaceutical Bulletin, 27 (6), 806-809.
- Nerya, O. et al. (2003). Glabrene and isoliquiritigenin as tyrosinase Inhibitors from licorice roots. Journal of Agriculture Food Chemistry, 51 (5), 1201-1207.
- No, J.K. et al. (1999). Inhibition of tyrosinase by green tea components. Pharmacology Letters, 65 (21), 241-246.
- Nohynek, G.J. et al. (2004). An assessment of the genotoxicity and human health risk of topical use of kojic acid (5-hydroxy-2-(hydroxymethyl)-4H-pyran-4-one). Food and Chemical Toxicology, 42 (1), 96-105.
- Okawa, M., Kinjo, J. & Nohara, T. (2001). DPPH (1,1-diphenyl-2-picrylhydrazyl) radical scavenging activity of flavonoids obtained from some medicinal plants. Biological and Pharmaceutical Bulletin, 24 (10), 1202-1205.
- Pinnell, S.R. (2003). Cutaneous photodamage, oxidative stress, and topical antioxidant protection. Journal of American Academic Dermatology, 48 (1), 1-19.
- Ponec, M. (2002). Skin constructs for replacement skin tissue for *in vitro* testing. Advanced Drug Delivery, 54 (1), 19-30.
- Roh, J.S. et al. (2004). Inhibitory effects of active compounds isolated from safflower (*Carthamus tinctorius* L.) seed for melanogenesis. Biological and Pharmaceutical Bulletin, 27(12), 1976-1978.
- Sanchez-Ferrer, A. et al. (1995). Tyrosinase: a comprehensive review of its mechanism. Biochemical et Biophysica Acta, 1247 (1), 1-11.
- Scott, G. (2002). Rac and Rho: the story behind melanocyte dendrite formation. Pigment Cell Research, 15 (5), 322-330.
- Seiberg, M. et al. (2000). The protease-activated receptor 2 regulates pigmentation via keratinocyte-melanocyte interactions. Cell Research, 254 (1), 25-32.

- Seiberg, M. (2001). Keratinocyte-melanocyte interactions during melanosome transfer. Pigment Cell Research, 14 (4), 236-242.
- Shimizu, K. et al. (1998). The inhibitory components from *Artocarpus incisus* on melanin biosynthesis. Planta Medica, 64 (5), 408-412.
- Shimizu, K., Kondo, R., & Sakai, K. (2002). The skin-lightening effects of artocarpin on UVB-induced pigmentation. Planta Medica, 68 (1), 79-81.
- Stanojevic, M. et al. (2004). Ultraviolet radiation and melanogenesis. Archive of Oncology, 12 (4), 203-205.
- Sugimoto, K. et al. (2004). Inhibitory effect of α -arbutin on melanin synthesis in cultured human melanoma cells and a three-dimensional human skin model. Biological and Pharmaceutical Bulletin, 27(4), 510-514.
- Taylor, S.C. (2002). Skin of color: biology, structure, function, and implications for dermatologic disease. American Academy of Dermatology, 46 (2), 41-62.
- Tiedtke, J., Morel, J. & Marks, O. (April 11, 2004). Depigmentation factor bioflavonoids – a safe and effective skin lightener based on encapsulated citrus bioflavonoids. Retrived on November 10, 2002 from: http://www.cosmetochem.ch/upload/docs/prodlist/depigmentation_F_B_Article.pdf.
- Virador, V.M. et al. (2002). Influence of melanocyte-stimulating hormone and of ultraviolet radiation on the transfer of melanosomes to keratinocytes. The FASEB Journal, 16 (1), 105-107.
- Walters, K.A (Ed). (2002). The structure and function of skin. New York: Marcel Dekker.
- Wulf, H.C. et al. (2004). Skin aging and natural photoprotection. Micron, 35(3), 185-191.