

LIST OF CONTENTS

Chapter	Page
I INTRODUCTION.....	1
The rationale of the study	1
Objectives of the study	3
Expected output of the study	4
II LITERATURE REVIEWS.....	5
<i>Artocarpus incisa</i> (Breadfruit).....	5
Skin structure.....	8
Epidermis.....	9
Dermis	12
Subcutaneous	12
Skin color.....	12
UV induced pigmentation.....	16
Depigmentation and skin whitening agents.....	17
Antioxidant properties as potent tyrosinase inhibitors.....	20
Post-inflammatory hyperpigmentation of skin.....	22
Nanoemulsions	23
Phase inversion temperature method (PIT)	24
Permeability pathway	25
<i>In vitro</i> evaluation of transdermal delivery	27
Evaluation of efficacy whitening products by scientific equipment....	29
III MATERIALS AND METHODS	31
Materials	31
Apparatus.....	33
Methods	34

LIST OF CONTENTS (CONT.)

Chapter	Page
IV RESULTS AND DISCUSSION	44
The characteristics of <i>A. incisus</i> extract	44
The content of artocarpin in <i>A. incisus</i> extract.....	45
<i>In vitro</i> efficacy of <i>A. incisus</i> extract on melanogenesis inhibition	48
<i>In vitro</i> antioxidant activity of <i>A. incisus</i> extract	52
<i>In vitro</i> anti-inflammatory activity of <i>A. incisus</i> extract.....	56
Physicochemical characterization of nanoemulsions	58
Stability studies of nanoemulsion containing <i>A. incisus</i> extract	63
<i>In vitro</i> permeation studies through mouse skin.....	66
<i>In vivo</i> studies of UVB-induced hyperpigmentation in C57BL/6 mice ..	68
V CONCLUSION	73
REFFERENCES	75
APPENDIXES.....	87
BIOGRAPHY	119

LIST OF TABLES

Table	Page
1 The appearance, percentage of yield and moisture content of the <i>A. incisus</i> 's heartwood extract	44
2 The conditions and moisture content of <i>A. incisus</i> 's heartwood extract	45
3 The percentage of free radical scavenging of <i>A. incisus</i> extract dissolved in DMSO	53
4 The percentage of free radical scavenging of L-ascorbic acid dissolved in deionized water	54
5 The percentage of free radical scavenging of butylated hydroxytoluene (BHT) dissolved in methanol	55
6 The inhibitory effect of <i>A. incisus</i> extract on TNF- α production and on cell viability of RAW264.7 cells stimulated by 1 μ g/mL of LPS (Data are in means \pm S.D. of triplicate study).....	57
7 The inhibitory effect of prednisolone on TNF- α production and on cell viability of RAW264.7 cells stimulated by 1 μ g/mL of LPS (Data are in means \pm S.D. of triplicate study).....	57
8 The peak areas of artocarpin at various concentrations of artocarpin standard regarding to the absorbance at wavelength of 282 nm.....	90
9 The content of artocarpin contained in extract of <i>A. incisus</i> 's heartwood..	91
10 The cell number of time course study	93
11 The effect of <i>A. incisus</i> extract on cells number	94
12 The melanin content	95
13 The antioxidant activity of <i>A. incisus</i> extract	96
14 The antioxidant activity of L-ascorbic acid	97
15 The antioxidant activity of butylated hydroxytoluene (BHT).....	99
16 The inhibitory effect of <i>A. incisus</i> extract on TNF- α release	102
17 The inhibitory effect of prednisolone on TNF- α release	104
18 The inhibitory effect of <i>A. incisus</i> extract on cell viability	106

LIST OF TABLES (CONT.)

Table	Page
19 The inhibitory effect of prednisolone on cell viability	108
20 Mean droplet size and polydispersity index value of nanoemulsions formulated with cetareth-10.....	110
21 Mean droplet size and polydispersity index value of nanoemulsions formulated with cetareth-20.....	111
22 Artocarpin remaining in nanoemulsion under 7 heat-cool cycles and after 1-3 months	112
23 Droplet size of nanoemulsion under 7 heat-cool cycles and after 1-3 months	113
24 Viscosity of nanoemulsion containing <i>A. incisus</i> extract under 7 heat- cool cycles and after 1-3 months	114
25 pH values of nanoemulsion containing <i>A. incisus</i> extract under 7 heat- cool cycles and after 1-3 months.....	115
26 The percentage of cumulative of artocarpin release of nanoemulsion were determined within 24 hr.....	116
27 The percentage of cumulative of artocarpin release of solution were determined within 24 hr.....	117
28 The Effect of nanoemulsion containing <i>A. incisus</i> extract and <i>A. incisus</i> extract solution on UVB-induced hyperpigmentation in C57BL/6 mice skin.....	118

LIST OF FIGURES

Figure	Page
1 <i>Artocarpus incisus</i>	5
2 Chemical structure of active compounds isolated from <i>A. incisus</i> heartwood extract.....	8
3 Skin structure	9
4 Epidermis structure	10
5 The melanocyte	13
6 The distribution of melanin in the epidermis	14
7 The melanin synthesis pathway.....	15
8 Chemical structures of whitening agents.....	18
9 Free radical form of DPPH (A) and reduced form of DPPH (B) after accepted an electron or hydrogen atom	21
10 The principle of the PIT method.....	25
11 Transcellular and intercellular route.....	26
12 Follicular route	26
13 Franz-type diffusion cell apparatus	28
14 (A) Mexameter and (B) Mexameter's probe (Model MX 18, Courage and Khazaka Electronic GmbH, Cologne, Germany).....	30
15 The diagram of UVB-induced hyperpigmentation study (A) back hair removal with shaving razor, (B) initial pigmentation before UV irradiation, (C) back skin irradiation from UVB lamps, (D) back skin with hyperpigmentation after UV irradiation.....	42
16 The appearance of <i>A. incisus</i> diethyl ether extract	45
17 Chromatogram of 0.2 mg/ml artocarpin standard	46
18 Chromatogram of 0.04 mg/ml artocarpin standard	46
19 Chromatogram of 1.0 mg/ml <i>A. incisus</i> ether extract	47

LIST OF FIGURES (CONT.)

Figure	Page
20 The calibration curve of artocarpin satandard	47
21 Growth curve of melanocyte B16F1 mouse melanoma cell line (Each point represents mean \pm S.D. of triplicate study).....	48
22 Morphology of melanocyte B16F1 melanoma cells of day 1 (A) and day 5 (B) at magnification of 20X	49
23 Inhibitory effects of <i>A. incisus</i> extract and kojic acid on melanin Synthesis in B16F1 cells (Data are expressed as percent of control, and each point represents mean \pm S.D. of triplicate study).....	50
24 Morphology of melanocyte B16F1 melanoma cells after being treated with 0.1% DMSO (control) and with 40 μ g/mL Kojic Acid for 4 days at magnification of 40X.....	51
25 Morphology of melanocytes B16F1 melanoma cells after being treated with 0.1% DMSO (control) and with 40 μ g/mL of <i>A. incisus</i> extract for 4 days at magnification of 40X	51
26 The effect of <i>A. incisus</i> on melanocyte B16F1 melanoma cells number (Each point represents mean \pm S.D. of triplicate study. Significant differences from control; ** p <0.01).....	52
27 Antioxidant activity of <i>A. incisus</i> extract	53
28 Antioxidant activity of L-ascorbic acid	54
29 Antioxidant activity of <i>A. incisus</i> ether extract, L-ascorbic acid and butylated hydroxytolue (BHT).....	55

LIST OF FIGURES (CONT.)

Figure	Page
30 Transmission electron photomicrograph of nanoemulsion formulation consisted of 0.02% w/w <i>A. incisus</i> extract, 41.62% w/w isopropyl myristate, 8% w/w cetareth-10, 5% w/w GMS and 0.03% w/w carbopol 940.....	58
31 Effect of cetareth-10 concentration at 4, 8, 10, 12 and 16% w/w while maintaining the amount of GMS at 5% w/w on the droplet size of nanoemulsions (Each bar represents mean±S.D. of triplicate study).....	60
32 Effect of GMS concentration varied from 1-8% w/w while maintaining the amount of cetareth-10 at 8% w/w on the droplet size of nanoemulsions (Each bar represents mean±S.D. of triplicate study).....	61
while maintaining the amount of GMS at 5% w/w on the droplet size of nanoemulsions (Each bar represents mean±S.D. of triplicate study).....	62
34 Effect of GMS concentration varied from 1-8% w/w while maintaining the amount of cetareth-20 at 8% w/w on the droplet size of nanoemulsions (Each bar represents mean±S.D. of triplicate study).....	63
35 Percentage of artocarpin remaining in nanoemulsion under normal condition for various periods and stress condition employing 7 heat cool cycles (Each bar represents mean±S.D. of triplicate study. Significant differences from freshly prepare; * $p < 0.05$)......	64
36 Changes in droplet size of nanoemulsion under normal condition for various periods and stress condition employing 7 heat cool cycles (Each bar represents mean±S.D. of triplicate study. Significant differences from initial; * $p < 0.05$)......	65

LIST OF FIGURES (CONT.)

Figure	Page
37 Changes in viscosity of nanoemulsion under normal condition for various periods and stress condition employing 7 heat-cool cycles (Each bar represents mean±S.D. of triplicate study. Significant differences from initial; * $p < 0.05$).....	65
38 Changes in pH of nanoemulsion under normal condition for various periods and stress condition employing 7 heat cool cycles (Each bar represents mean±S.D. of triplicate study).....	66
39 <i>In vitro</i> release profile of artocapin from nanoemulsion and <i>A. incisus</i> extract solution (Each point represents mean±S.D. of triplicate study. Significant differences from control; ** $p < 0.01$).....	67
40 Effect of nanoemulsion containing <i>A. incisus</i> extract and <i>A. incisus</i> solution on UVB-induced hyperpigmentation in C57BL/6 mouse skin; all photographs are from the same animal (A) initial pigment after UV irradiation and after the topical application, (B) 4 weeks, (C) 6 weeks.....	69
41 Changes of M-value after the daily topical applications (A) and the degree of pigmentation (Δ M-value) before and 6 weeks after daily topical application (B) of the nanoemulsion containing <i>A. incisus</i> extract and <i>A. incisus</i> extract solution. (Each point represents mean±S.D. of triplicate study. Significant differences from control; * $p < 0.05$).....	70
42 The HPLC chromatogram of artocarpin standard of 0.2 mg/mL	88
43 The HPLC chromatogram of artocarpin standard of 0.04 mg/mL.....	88
44 The HPLC chromatogram of <i>A. incisus</i> extract.....	89
45 The calibration curve of artocarpin standard.....	91
46 The standard curve of mouse TNF- α ELISA Ready-SET-Go.....	101